

Algebraic Points of Any Given Degree on the Affine Equation Curve $y^{11} = x^4(x - 1)^4$

Mouhamadou Diaby Gassama, Chérif Mamina Coly, Oumar Sall

Mathematics and Applications Laboratory, Faculty of Science and Technology, Assane Seck University in Ziguinchor, Ziguinchor, Senegal

Email address :

m.gassama1299@zig.univ.sn (Mouhamadou Diaby Gassama), osall@univ-zig.sn (Oumar Sall), c.coly1309@zig.univ.sn (Chérif Mamina Coly)

To cite this article :

Mouhamadou Diaby Gassama, Chérif Mamina Coly, Oumar Sall. Algebraic Points of Any Given Degree on the Affine Equation Curve $y^{11} = x^4(x - 1)^4$. *American Journal of Applied Mathematics*. Vol. 11, No. 5, 2023, pp. 89-94. doi : 10.11648/j.ajam.20231105.12

Received : September 29, 2022 ; **Accepted :** July 24, 2023 ; **Published :** November 1, 2023

Abstract : The quotients of Fermat curves $C_{r,s}(p)$ are studied by SALL who extends the work of Gross and Rohlich. Among these studies are the cases $C_{r,s}(11)$ for $r = s = 1$. COLY and Sall have explicitly determined the algebraic points of degree at most 3 on \mathbb{Q} for the cases $C_{r,s}(11)$ for $r = s = 2$. Our work focuses we determine explicitly the algebraic points of a given degree over on the curve $C_{4,4}(11)$ of affine equation $y^{11} = x^4(x - 1)^4$ which is a special case of Fermat quotient curves. Our study concerns the cases $C_{r,s}(11)$ for $r = s = 4$. This note completes previous work of Gassama and Sall who explicitly determined the algebraic points of degree at most three on the even curve. It seems that the finiteness of the Mordell-Weil group of rational points of the Jacobien $J_{4,4}(11)(\mathbb{Q})$ is an essential condition. So to determine the algebraic points on the curve $C_{4,4}(11)$ we need a finiteness of the Mordell-Weil group of rational points of the Jacobien $J_{4,4}(11)(\mathbb{Q})$. The Mordell-Weil group $J_{4,4}(11)(\mathbb{Q})$ of rational points of the Jacobien is finite according to Faddev. Our note is in this framework. Our essential tools in this note are the Mordell-Weil group $J_{4,4}(11)(\mathbb{Q})$ of the Jacobien of $C_{4,4}(11)$ the Abel-Jacobi theorem and the study of linear systems on the curve $C_{4,4}(11)$. The result obtained concerns some quotients of Fermat curves. Indeed, the curve of affine equation $y^{11} = x^4(x - 1)^4$, we made an extension of the work of Gassama and Sall by explicitly determining the algebraic points of given degree on the curve $C_{4,4}(11)$ and this is what makes this note very interesting.

Keywords : Mordell-Weil Group, Jacobian, Galois Conjugates

1. Introduction

Let \mathcal{C} be a smooth algebraic curve defined on \mathbb{Q} . Let K be a field of numbers we note $\mathcal{C}(K)$ the set of points on \mathcal{C} with coordinates in K , and $\bigcup_{[K:\mathbb{Q}] \leq d} \mathcal{C}(K)$ the set of points on \mathcal{C} with coordinates in K of degree at most d on \mathbb{Q} . The degree of a point R of \mathcal{C} algebraic on \mathbb{Q} is defined as the degree of its defining field on \mathbb{Q} ; in other words $\deg(R) = [\mathbb{Q}(R) : \mathbb{Q}]$.

When \mathcal{C} is of genus $g \geq 2$, we know since Faltings [5] that the set of rational points $\mathcal{C}(\mathbb{Q})$ is finite. A generalization of this theorem to subvarieties of abelian varieties obtained by Vojta and Faltings [4, 16] can also be used to qualitatively describe $\bigcup_{[K:\mathbb{Q}] \leq d} \mathcal{C}(K)$.

In this note, our work will consist in the study of some particular cases, where particular cases, where we can determine explicitly the algebraic points of any degree on the curve of affine equation $C_{4,4}(11)$: $y^{11} = x^4(x - 1)^4$. \mathcal{C} corresponds to the curve $C_{4,4}(11)$.

Our curve $C_{4,4}(11)$ with affine equation $y^{11} = x^4(x - 1)^4$ is a special case of the quotients of Fermat curves $C_{r,s}(p)$: $y^p = x^r(x - 1)^s$, $1 \leq r, s$; $r + s \leq p - 1$ studied in [10 - 12].

The cases $C_{r,s}(11)$ for $r = s = 2$ are studied in [1]. See [2, 3, 5, 8, 13] for other explicit examples. Indeed, \mathcal{C} corresponds to the curve $C_{4,4}(11)$. The curves $C_{r,s}(p)$ are quotients of F_p [7, 16].

We denote by $J_{4,4}(11)$ the Jacobien of $C_{4,4}(11)$ and by $j(P)$ the class denoted $[P - P_\infty]$ of $P - P_\infty$, that is to say j is the Jacobien fold $C_{4,4}(11) \rightarrow J_{4,4}(11)$. The Mordell-Weil group $J_{4,4}(11)(\mathbb{Q})$ of the rational points of the Jacobien is finite [6, 7, 14]. The curve $C_{4,4}(11)$ in projective is $C_{4,4}(11) : Y^{11} = X^4 Z^7 (X - Z)^4$. Let us note P_0 , P_1 and P_∞ the points defined by : $P_0 = (0, 0, 1)$; $P_1 = (1, 0, 1)$ and $P_\infty = (1, 0, 0)$.

Gassama and Sall [9], determined the set of algebraic points

of degree at most 3 on \mathbb{Q} of the curve with affine equation $\mathcal{C}_{4,4}(11)$.
 $y^{11} = x^4(x - 1)^4$. We extend this result, giving an explicit description of the points of any degree on \mathbb{Q} on the curve

In this note we determine the set :

$$\bigcup_{[K : \mathbb{Q}] \leq l} \mathcal{C}_{4,4}(11)(K)$$

Our main result is the following :

Théorème 1. The set of algebraic points of at most any degree l on \mathbb{Q} on the curve $\mathcal{C}_{4,4}(11)$ is given by :

$$\bigcup_{[K : \mathbb{Q}] \leq l} \mathcal{C}_{4,4}(11)(\mathbb{K}) = \mathcal{G}_0 \cup \left(\bigcup_{1 \leq n \leq 10} \mathcal{G}_n \right)$$

where :

$$\mathcal{G}_0 = \left\{ \left(\left(\begin{array}{c} \sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{2}} \\ -\frac{\sum_{i=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}}} \end{array} \right), y \right) \mid \begin{array}{l} (\alpha_0 \wedge \beta_0) \neq 0, \alpha_{\frac{l}{2}} \neq 0 \text{ if } l \text{ is even, } \beta_{\frac{l-11}{2}} \neq 0 \text{ if } l \text{ is odd and} \\ y \text{ solution of the equation} \\ y^{11} \left(\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}} \right)^4 = \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{2}} \right)^2 \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{2}} + \sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}} \right)^2 \end{array} \right\}$$

$$\mathcal{G}_n = \left\{ \left(\left(\begin{array}{c} \sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{2}} \\ -\frac{\sum_{i=0}^{\frac{l-11}{2}} \alpha_i y^{\frac{i}{2}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}}} \end{array} \right), y \right) \mid \begin{array}{l} \beta_0 \neq 0, \alpha_{\frac{l+11-n}{2}} \neq 0 \text{ if } l \text{ is even, } \beta_{\frac{l-n}{2}} \neq 0 \text{ if } l \text{ is odd and} \\ y \text{ solution of the equation} \\ y^n \left(\sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^4 = \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha_i y^{\frac{i+n-11}{2}} \right)^2 \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha_i y^{\frac{i}{2}} + \sum_{j=0}^{\frac{l-n}{2}} \beta_j y^{\frac{j}{2}} \right)^2 \end{array} \right\}$$

2. Auxiliary Results

For a divisor D on C , we denote $\mathcal{L}(D)$ the $\bar{\mathbb{Q}}$ -vector space of rational functions f defined on \mathbb{Q} such that $f = 0$ or $\text{div}(f) \geq -D$; $l(D)$ denotes the $\bar{\mathbb{Q}}$ -dimension of $\mathcal{L}(D)$.

Lemma 1. we have : $\mathcal{J}_{4,4}(11)(\mathbb{Q}) \cong \mathbb{Z}/11\mathbb{Z}$

Demonstration : According Gross and Rohrlich ([7], page. 219), we have : $\mathcal{J}_{4,4}(11)(\mathbb{Q})_{\text{torsion}} \cong \mathbb{Z}/11\mathbb{Z}$ and According to Faddeev [6], we have : $\mathcal{J}_{4,4}(11)(\mathbb{Q})_{\text{torsion}} \cong \mathcal{J}_{4,4}(11)(\mathbb{Q})$.

Lemma 2. For the curve $\mathcal{C}_{4,4}(11) : y^{11} = x^4(x - 1)^4$, we have :

1. $\text{div}(x) = 11P_0 - 11P_\infty$,
2. $\text{div}(x - 1) = 11P_1 - 11P_\infty$,
3. $\text{div}(y) = 4P_0 + 4P_1 - 8P_\infty$.

Demonstration : (see [11], Lemme 1)

Corollary 1. The following results are the consequences of Lemma 2.

1. $4j(P_0) = -4j(P_1)$,
2. $11j(P_0) = 11j(P_1) = 0$

So $j(P_0)$ and $j(P_1)$ generate the same subgroup $\mathcal{J}_{4,4}(11)(\mathbb{Q})$ isomorphic to $\mathbb{Z}/11\mathbb{Z}$. Thus we have :

$$\mathcal{J}_{4,4}(11)(\mathbb{Q}) \cong \mathbb{Z}/11\mathbb{Z} = \{nj(P_0), 0 \leq n \leq 10\}.$$

Lemma 3. A \mathbb{Q} -base of $\mathcal{L}(mP_\infty)$ is given by :

$$\mathfrak{B}_m = \left\{ \left(\frac{y^3}{x(x - 1)} \right)^i, 0 \leq i \leq \frac{m}{2} \right\} \cup \left\{ x \left(\left(\frac{y^3}{x(x - 1)} \right)^j \right), 0 \leq j \leq \frac{m-11}{2} \right\}$$

Demonstration :

It is easy to show that \mathcal{B}_m is a free family, it then remains to show that $\text{card } \mathcal{B}_m = \dim \mathcal{L}(mP_\infty)$. We know that the genus of \mathcal{C} is $g = \frac{11-1}{2} = 5$. Since the curve has genus 5, by the Riemann-Roch theorem, we have $\dim \mathcal{L}(mP_\infty) = m - g + 1 = m - 4$ as soon as $m \geq 2g - 1 = 9$. For $m < 9$, $\dim \mathcal{L}(mP_\infty)$ is given by Clifford's theorem, which says that $\dim \mathcal{L}(mP_\infty) \geq \frac{1}{2} \deg(mP_\infty) + 1 = \frac{1}{2}m + 1$. Two cases are possible :

1. 1st case : suppose that m is even, then $m = 2h$, we obtain :

$$i \leq \frac{m}{2} \Leftrightarrow i \leq \frac{2h}{2} = h \text{ in the same way } j \leq \frac{m-11}{2} \Leftrightarrow j \leq \frac{2h-11}{2} \Leftrightarrow j \leq h - \frac{11}{2} \\ \Rightarrow j < h - \frac{10}{2} = h - 5 \Rightarrow j \leq h - 6. \text{ So we have :}$$

$$\mathcal{B}_m = \left\{ 1, \left(\frac{y^3}{x(x-1)} \right), \dots, \left(\frac{y^3}{x(x-1)} \right)^h \right\} \cup \left\{ x, x \left(\frac{y^3}{x(x-1)} \right), \dots, x \left(\frac{y^3}{x(x-1)} \right)^{h-6} \right\}.$$

We deduce that : $\text{card } \mathcal{B}_m = h + 1 + h - 6 + 1 = 2h - 4 = m - 4 = \dim \mathcal{L}(mP_\infty)$.

2. 2nd case : suppose that m is odd, then $m = 2h + 1$, we obtain :

$$i \leq \frac{m}{2} \Leftrightarrow i \leq \frac{2h+1}{2} \Leftrightarrow i \leq h + \frac{1}{2} \Rightarrow i < h + 1 \Rightarrow i \leq h \text{ in the same way} \\ j \leq \frac{m-11}{2} \Leftrightarrow j \leq \frac{2h-10}{2} = h - 5. \text{ So we have :}$$

$$\mathcal{B}_m = \left\{ 1, \left(\frac{y^3}{x(x-1)} \right), \dots, \left(\frac{y^3}{x(x-1)} \right)^h \right\} \cup \left\{ x, x \left(\frac{y^3}{x(x-1)} \right), \dots, x \left(\frac{y^3}{x(x-1)} \right)^{h-6} \right\}.$$

We deduce that : $\text{card } \mathcal{B}_m = h + 1 + h - 5 + 1 = 2h - 3 = m - 4 = \dim \mathcal{L}(mP_\infty)$.

3. Demonstration of the Theorem

Let $R \in \mathcal{C}_{4,4}(11)(\bar{\mathbb{Q}})$ with $[\mathbb{Q}(R) : \mathbb{Q}] = l$. Notons R_1, \dots, R_l Let R and l be the conjugates of R , and work with $t = [R_1 + \dots + R_l - lP_\infty]$ which is a point of $\mathcal{J}_{4,4}(11)(\mathbb{Q}) = \{nj(P_0), 0 \leq n \leq 10\}$; donc $t = nj(P_0)$ with $0 \leq n \leq 10$, ains

$$[R_1 + \dots + R_l - lP_\infty] = nj(P_0) \text{ with } 0 \leq n \leq 10 \quad (i)$$

We discuss according to the values of $n \in \{0, \dots, 10\}$:

1. case : $n = 0$. Th formula (i) becomes : $[R_1 + \dots + R_l - lP_\infty] = 0$.

According to the Abel-Jacobi theorem there exists then a rational function f defined on \mathbb{Q} such that $\text{div}(f) = R_1 + \dots + R_l - lP_\infty$, donc $f \in \mathcal{L}(lP_\infty)$.

According to Lemma 3, we have, :

$$f = \sum_{i=0}^{\frac{l}{2}} a_i \left(\frac{y^3}{x(x-1)} \right)^i + \sum_{j=0}^{\frac{l-11}{2}} b_j x \left(\frac{y^3}{x(x-1)} \right)^j$$

with $a_0 \neq 0$ (otherwise one of the R_i should be equal to P_0 , which would be absurd), $a_{\frac{l}{2}} \neq 0$ if l is even (otherwise one of R_i would be equal to P_∞ , which would be absurd) and $b_{\frac{l-11}{2}} \neq 0$ if l is odd (otherwise one of R_i would be equal to P_∞ , which would be absurd).

At the points R_i , we have :

$$\sum_{i=0}^{\frac{l}{2}} a_i \left(\frac{y^3}{x(x-1)} \right)^i + \sum_{j=0}^{\frac{l-11}{2}} b_j x \left(\frac{y^3}{x(x-1)} \right)^j = 0$$

$$\text{Hence } x = - \frac{\sum_{i=0}^{\frac{l}{2}} a_i \left(\frac{y^3}{x(x-1)} \right)^i}{\sum_{j=0}^{\frac{l-11}{2}} b_j \left(\frac{y^3}{x(x-1)} \right)^j}.$$

The relation $y^{11} = x^4(x - 1)^4$, involves $\frac{y^3}{x(x-1)} = \pm y^{\frac{1}{4}}$. Thus we have :

$$x = -\frac{\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}}} \text{ where } \begin{cases} \alpha_i = \pm a_i \\ \beta_j = \pm b_j \end{cases}.$$

And as a result :

$$\begin{aligned} y^{11} = x^4(x - 1)^4 &\iff y^{11} = \left(-\frac{\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}}} \right)^4 \left(-\frac{\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}}} - 1 \right)^4 \\ &\iff y^{11} \left(\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^8 = \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} \right)^4 \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} + \sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^4 \end{aligned}$$

The expression :

$$y^{11} \left(\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^8 = \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} \right)^4 \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} + \sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^4$$

is an equation of degree l in y ; indeed

The first member of the equation is of degree at most equal to $11 + 8 \left(\frac{\frac{l-11}{2}}{4} \right) = l$ and the second member of the equation is of degree equal to $2 \times 4 \left(\frac{\frac{l}{2}}{4} \right) = l$.

We thus obtain a family of points of degree l :

$$\mathcal{G}_0 = \left\{ \left(-\frac{\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}}}{\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}}}, y \right) \mid \begin{array}{l} \alpha_0 \neq 0, \alpha_{\frac{l}{2}} \neq 0 \text{ if } l \text{ is even, } \beta_{\frac{l-11}{2}} \neq 0 \text{ if } l \text{ is odd and} \\ y \text{ solution of the equation} \\ y^{11} \left(\sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^8 = \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} \right)^4 \left(\sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{4}} + \sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{4}} \right)^4 \end{array} \right\}$$

2. case : $n \in \{1, \dots, 10\}$. Formula (i) becomes : $[R_1 + \dots + R_l - lP_\infty] = nj(P_0)$, we deduce that $[R_1 + \dots + R_l + (11 - n)P_0 - (l + 11 - n)P_\infty] = 0$.

According to the Abel-Jacobi theorem, there exists then a rational function f defined on \mathbb{Q} such that :

$$\text{div}(f) = R_1 + \dots + R_l + (11 - n)P_0 - (l + 11 - n)P_\infty,$$

so $f \in \mathcal{L}((l + 11 - n)P_\infty)$.

According to Lemma 3, we have :

$$f = \sum_{i=0}^{\frac{l+11-n}{2}} a_i \left(\frac{y^6}{x(x-1)} \right)^i + \sum_{j=0}^{\frac{l-n}{2}} b_j x \left(\frac{y^6}{x(x-1)} \right)^j$$

with $b_0 \neq 0$ (otherwise one of R_i should be equal to P_0 , which would be absurd), $a_{\frac{l}{2}} \neq 0$ if l is even (otherwise one of R_i would be equal to P_∞ , which would be absurd) and $b_{\frac{l-n}{2}} \neq 0$ if l is odd (otherwise one of R_i would be equal to P_∞ , which would be absurd).

At points R_i , we have :

$$\sum_{i=11-n}^{\frac{l+11-n}{2}} a_i \left(\frac{y^6}{x(x-1)} \right)^i + \sum_{j=0}^{\frac{l-n}{2}} b_j x \left(\frac{y^6}{x(x-1)} \right)^j = 0$$

$$\text{Hence } x = - \frac{\sum_{i=11-n}^{\frac{l+11-n}{2}} a_i \left(\frac{y^6}{x(x-1)} \right)^i}{\sum_{j=0}^{\frac{l-n}{2}} b_j \left(\frac{y^6}{x(x-1)} \right)^j}.$$

In addition $\frac{y^6}{x(x-1)} = \pm y^{\frac{1}{2}}$, we deduce that :

$$x = - \frac{\sum_{i=0}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i}{2}}}{\sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}}} \text{ where } \begin{cases} \alpha'_i = \pm a_i \\ \beta'_j = \pm b_j \end{cases}$$

From the relation $y^{11} = x^2(x-1)^2$, we deduce, using the same procedure as the previous case, the following equation in y :

$$y^{11} \left(\sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^4 = \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i}{2}} \right)^2 \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i}{2}} + \sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^2$$

This equation becomes :

$$y^n \left(\sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^4 = \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i+n-11}{2}} \right)^2 \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i}{2}} + \sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^2$$

which is an equation of degree l . Indeed :

For l is even or odd, the first member of the equation is of degree equal to $n + 4 \left(\frac{\frac{l-n}{2}}{2} \right) = l$ and the second member of the equation is of degree equal to $2 \times \left(\frac{\frac{l+11-n}{2} + n - 11}{2} \right) + 2 \times \left(\frac{\frac{l+11-n}{2}}{2} \right) = l$.

We thus obtain a family of points of degree l :

$$\mathcal{G}_n = \left\{ \left(\begin{pmatrix} \sum_{i=0}^{\frac{l}{2}} \alpha_i y^{\frac{i}{2}} \\ -\frac{i=0}{\frac{l-11}{2}} \\ \sum_{j=0}^{\frac{l-11}{2}} \beta_j y^{\frac{j}{2}} \end{pmatrix}, y \right) \mid \begin{array}{l} \beta_0 \neq 0, \alpha_{\frac{l+11-n}{2}} \neq 0 \text{ si } l \text{ est pair, } \beta_{\frac{l-n}{2}} \neq 0 \text{ if } l \text{ is odd and} \\ y \text{ solution of the equation} \\ y^n \left(\sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^4 = \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i+n-11}{2}} \right)^2 \left(\sum_{i=11-n}^{\frac{l+11-n}{2}} \alpha'_i y^{\frac{i}{2}} + \sum_{j=0}^{\frac{l-n}{2}} \beta'_j y^{\frac{j}{2}} \right)^2 \end{array} \right\}$$

4. Conclusion

Our note focuses on the determination of algebraic points on the curve $\mathcal{C}_{4,4}(11)$ of affine equation $y^{11} = x^4(x-1)^4$. The curve $\mathcal{C}_{4,4}(11)$ is a special case of the quotients of Fermat curves.

We have determined the algebraic points of any given degree on the given curve. To be able to do it one of the bases was determined and then the determination of given degree 1 of algebraic points. The determination of the algebraic points of degree exactly 1 on the given curve remains to be studied.

References

- [1] C. M. Coly, O. Sall, points algébriques de degré au-plus 3 sur la courbe d'équation affine $y^{11} = x^2(x - 1)^2$, Annales Mathématiques Africaines Volume 8 (2020) pp. 27-32.
- [2] O. Debarre, R. Fahlouai, Abelian varieties and curves in $W(C)$ and points of bounded degree on algebraic curves, Compositio Math. 88 (1993) 235-249.
- [3] O. Debarre, M. Klassen, Points of low degree on smooth plane curves, J. Reine Angew. Math. 446 (1994) 81-87.
- [4] Diophantine approximation on Abelian varieties, Ann. Math. 133 (1991) 549-576.
- [5] D. Faddeev, On the divisor class groups of some algebraic curves, Dokl. Akad. Nauk SSSR 136 (1961) 296-298. English translation : Soviet Math. Dokl. 2 (1) (1961) 67-69.
- [6] G. Faltings, Endlichkeitsätze für abelsche Varietäten über Zahlkörpern, Invent. Math. 73 (1983) 349-366.
- [7] B. Gross, D. Rohrlich, Some results on the Mordell-Weil group of the Jacobian of the Fermat curve, Invent. Math. 44 (1978) 201-224.
- [8] M. Klassen, P. Tzermias, Algebraic points of low degree on the Fermat quintic, Acta Arith. 82 (4) (1997) 393-401.
- [9] Mouhamadou Diaby Gassama, Oumar Sall. Algebraic Points of Degree at Most 3 on the Affine Equation Curve $y^{11} = x^4(x - 1)^4$. American Journal of Applied Mathematics. Vol. 10, No. 4, 2022, pp. 160-175. doi : 10.11648/j.ajam.20221004.15.
- [10] O. Sall, Points algébriques de petit degré sur les courbes de Fermat, C. R. Acad. Sci. Paris Sér. I 330 (2000) 67-70.
- [11] O. Sall, Points cubiques sur la quartique de Klein, C. R. Acad. Sci. Paris Sér. I 333 (2001) 931-934.
- [12] O. Sall, algebraic points on some Fermat curves and some quotients of Fermat curves : Progress, African Journal of Mathematical Physics Volume 8 (2010) 79-83.
- [13] O. Sall, M. Fall, points algébriques de petits degrés sur les courbes d'équations affines $y^{3n} = x(x-1)(x-2)(x-3)$, Annales Mathématiques Africaines (2015).
- [14] J. P. Serre, Lecture of Mordell weil theorem. the translated from the french and edited by martin brows from notes by michel Waldschmidt. Aspects of mathematics, E15 (1989).
- [15] P. Tzermias, Algebraic points of low degree on the Fermat curve of degree seven, Manuscript Math. 97 (4) (1998) 483-488.
- [16] P. Tzermias, Torsion parts of Mordell-Weil groups of Fermat Jacobians, Internat. Math. Res. Notices 7 (1998) 359-369.
- [17] Siegel's theorem in the compact case, Ann. Math. 133 (1991) 509-548.