

Pathophysiological Effects of Alcohol and Tobacco Consumption on Semen Parameters of Men Attending a Fertility Clinic in West Africa

Philip Vatcanarat Baysah¹, Aloysius Obinna Ikwuka^{1,*}, Francis Chigozie Udeh¹, Darius Popo Bleh², Thamara Vilorio³

¹College of Medicine and Health Sciences, American International University West Africa, Banjul, The Gambia

²Department of Laboratory Services, Medlink Clinic, Monrovia, Liberia

³College of Clinical Embryology and Artificial Reproduction Technology, University of Valencia, Valencia, Spain

Email address:

aloyssiussweet@yahoo.com (Aloysius Obinna Ikwuka)

*Corresponding author

To cite this article:

Philip Vatcanarat Baysah, Aloysius Obinna Ikwuka, Francis Chigozie Udeh, Darius Popo Bleh, Thamara Vilorio. Pathophysiological Effects of Alcohol and Tobacco Consumption on Semen Parameters of Men Attending a Fertility Clinic in West Africa. *American Journal of Biomedical and Life Sciences*. Vol. 11, No. 4, 2023, pp. 73-81. doi: 10.11648/j.ajbls.20231104.13

Received: July 12, 2023; Accepted: July 28, 2023; Published: August 5, 2023

Abstract: *Background:* Alcohol and tobacco consumption are prevalent global habits. There is existing evidence linking these habits to male infertility, although the impact they have on male fertility and reproductive outcomes are yet to be exhaustively investigated. *Aim:* This study aimed to study the pathophysiological effects of alcohol and tobacco consumption on semen parameters of men attending a fertility clinic in West Africa. *Materials and Methods:* Semen samples were collected from 196 men who attend Medlink Clinic, Monrovia, Liberia and parameters including sperm volume, count, motility, and morphology were evaluated according to the WHO criteria. The Chi-square test (χ^2) of the SPSS version 23 statistical software was used to test for hypotheses at a significance level of $p < 0.05$. *Results:* The mean age of the men was 37.8 ± 10.3 years. 20.9% of the study participants abstain from alcohol and tobacco, 38.8% consume alcohol, and 40.3% consume tobacco. Among the study participants, 36 (18.4%) had normospermia, 74 (37.8%) had oligospermia, 8 (4.1%) had cryptozoospermia, 5 (2.6%) had azoospermia, 58 (29.6%) had asthenozoospermia, and 15 (7.7%) had teratozoospermia. However, 78.9% frequent alcohol consumers, 47.4% non-frequent alcohol consumers, 77.8% non-alcohol consumers, 89.8% tobacco smokers, and 69.6% non-tobacco smokers had poor semen quality. The hypotheses tested proved that there is a significant relationship ($p = 0.024$) between alcohol/tobacco consumption and poor semen quality, whereas there is no significant relationship ($p = 0.56$) between non-consumption of alcohol/tobacco and good semen quality. There is also a significant relationship ($p = 0.0086$) between alcohol consumption only and poor semen quality. *Conclusion:* Alcohol and tobacco consumption reduce male sperm quality, but abstinence from both does not guarantee optimum semen quality. Alcohol and/or tobacco consumption adversely affect the semen parameters (sperm volume, count, motility, and morphology).

Keywords: Pathophysiology, Tobacco, Alcohol Consumption, Semen Parameters, Semen Quality, Male Fertility

1. Introduction

Gametes are ova and sperm cells which are haploid and have one copy of each type of chromosome i.e. 1–22 X or 1–22 Y [34]. Spermatogenesis is the process of sperm production, occurring in the seminiferous tubules of the testes. It takes approximately 74 days for a spermatogonium to mature into a

spermatozoon, with over 100 million spermatozoa produced daily [55]. Male fertility primarily relies on semen quality, encompassing parameters such as sperm volume, count, motility, and morphology [52].

Several factors including hazards linked to certain occupations (long distance driving, military, etc); previous history of genital infections (e.g. gonorrhea, orchitis, etc); previous history of surgery in the genital tract or inguinal

region; lifestyle choices (sedentary lifestyle, lack of exercise, sleep deprivation, etc); environmental factors (eg. exposure to ionizing radiation, pesticides, heat from tight underwears and hot baths) can adversely affect semen parameters and lead to male infertility. Infertility is defined as the inability to achieve a clinical pregnancy after one year of regular unprotected sexual intercourse – regular sexual intercourse is 2-3 times sexual intercourse per week [80]. Infertility affects 15-30% of couples of reproductive age [55, 56, 74]. Male infertility rates surpass those of females [41], causing significant emotional distress for affected couples [78].

Alcohol and tobacco consumption are prevalent worldwide [76, 77], and are known for their addictive nature and associated health risks [8, 9, 52]. Tobacco use has been reported as teratogenic, while excessive use of both substances can be lethal [40, 63, 64].

Moderate alcohol consumption (4–7 units/week) has shown a positive correlation with sperm quality [54]. However, prolonged abuse of alcohol and tobacco has been linked to male infertility, as they negatively impact spermatogenesis through various mechanisms [4, 11, 18, 43, 59]. Firstly, alcohol consumption has been reported to be directly toxic to the Leydig and Sertoli cells in the testes. Secondly, it has also been linked with anomalies in the metabolism of testosterone [19]. Thirdly, the morphology of spermatozoa and sperm counts have also been affected by the daily use of alcohol among men [50]. Outcomes such as coil-tailed spermatozoa, immature testicular cells, spermatozoa head breakage, and distention of the mid-piece have been reported as morphological alterations [50].

Numerous studies have confirmed that tobacco is generally harmful as it contains alkaloids and other dangerous components like Tobacco-Specific Nitrosamines (TSNAs), Propionic acid, 2- Naphylamine, Choline, Tabacinine, Tabacine, Mercury (Hg), Cadmium (Cd), N-nitrosornicotine, Tar, Pyrene, Nicotelline, Carbon monoxide and Cembrene [10, 40, 45, 63, 64, 77]. Of these alkaloids, nicotine accounts for 96-98% [10, 40], and causes high tobacco addiction, and has the most immediate pharmacological action [20]. The effects of tobacco on male fertility have been studied extensively but there is a controversy on the outcome. Some researchers identified significant adverse effects on sperm quality, while others like Halmenschlager, *et al* did not [17]. Cadmium (Cd) affects male fertility and endocrine functions by significantly reducing semen [51]. Moreover, cigarette smoking can elevate inflammatory reactions which cause leukocytosis in the testes [43, 48]. Major free radicals that are of physiological significance are superoxide anion, hydroxyl radical, and hydroperoxyl radical, while non-radical is hydrogen peroxide [33]. Cigarette smoking can also cause sperm DNA fragmentation, axonemal damage in the flagellum, and low sperm count [5, 18].

In addition, the interplay, role and effects of metabolic syndrome diseases on male fertility are still being investigated by different researchers. Metabolic syndrome diseases, MSD (Hypertension, Adiposity, Diabetes mellitus and Dyslipidemia)

are interrelated diseases with very high morbidity and mortality rates [21, 22, 24, 38, 65]. Results from different studies have shown that high levels of blood pressure, glucose and lipid metabolic disorders, asymptomatic hyperuricemia, activation of systemic immune inflammation and fibrogenesis, contribute to kidney damage [25-27, 29-31, 35, 37, 39, 66-69, 73]. Adiposity, diabetes mellitus and dyslipidemia have also been linked with erectile dysfunction.

Metabolic Syndrome Diseases require new and effective treatment options. Dapagliflozin which is a Sodium-Glucose Linked Transporter 2 (SGLT-2) inhibitor and Liraglutide which is a Glucagon-like Peptide 1 Receptor Agonist (GLP-1 RA) have been found to increase the effectiveness of treatment and improve the clinical course of type 2 diabetes mellitus and hypertension in patients with such comorbidities [23, 28, 32, 36, 70-72].

The record presented in 2018, showed that the total alcohol consumption per capita (per person) among Liberian males who are ≥ 15 years is 6.12 liters i.e. 6.12 liters and above of pure alcohol consumed over a calendar year [61]. In 2020, tobacco smoking among Liberian men was 14.3%, a decline from 15.1% in 2018, and also a decline from 16.6% in 2015 according to the World Bank data [60]. The 15.1% rate in 2018 ranked Liberia at 134 among different countries of the world. This shows that the civil awareness made and the necessary laws passed by the 52nd Legislation banning public smoking, have contributed to the decline in tobacco smoking. With these rates of alcohol and tobacco consumption among Liberian men, it is therefore necessary to examine their impact on male fertility by evaluating semen parameters.

2. Materials and Methods

2.1. Study Area

This study is a cross-sectional study conducted from June 2020 to July 2021, at Medlink Clinic, Monrovia, Liberia, where fertility experts and clinical embryologists carry out some levels of fertility tests and treatments in patients. This clinic is a multi-faceted health institution that collaborates with other government and private agencies for the analysis of samples for fertility, and the collection of samples for DNA analysis in other tertiary healthcare facilities.

2.2. Study Population

The study population is made of an estimated four hundred (400) men who visited the clinic within a period of fourteen (14) months.

2.3. Sample Size and Sampling Technique

Krejcie and Morgan's formula was used to estimate the sample size of this study.

Below is the formula:

$$n = \frac{X^2 N p(1-p)}{e^2 (N-1) + X^2 p(1-p)}$$

Where:

n is the sample size;

N is the population size (400);

e is the acceptable sampling error (5% or 0.05);

X^2 is the Chi-square of the degree of freedom 1 and confidence 95% (3.841);

p is the proportion of population (if unknown, 0.5)

$$n = \frac{(3.841) \cdot (400) \cdot (0.5[1-0.5])}{0.05^2 \cdot (400-1) + (3.841 \cdot 0.5[1-0.5])}$$

$$n = \frac{(3.841) \cdot (400) \cdot (0.5[1-0.5])}{0.05^2 \cdot (400-1) + (3.841 \cdot 0.5[1-0.5])}$$

$$n = 196$$

$$\text{Sampling Interval} = \frac{N}{n}$$

N = population size (400); n = sample size (196)

$$SI = \frac{400}{196}$$

$$SI \sim 2$$

The sampling technique used was a systematic sampling technique with a sampling interval of 2, selecting a sample size of 196 from the study population of 400 men visiting the clinic during the 14 months of this study. The participants included frequent alcohol consumers, non-frequent alcohol consumers, non-alcohol consumers, tobacco smokers, and non-smokers who met the inclusion criteria. Men who had undergone surgeries on their reproductive organs, men on long-term use of fertility medication, men with congenital anomalies like varicocele, hydrocele, hypospadias, epispadias, cryptorchidism, Klinefelter syndrome, men with mumps and evident orchitis, men over 65 years of age, and men who quit alcohol and tobacco usage within three months prior to this study, were excluded. Other patients who met the inclusion criteria were requested to participate, and voluntarily consented to participate after making informed decision.

2.4. Data Collection Method

Semen samples were obtained from patients enrolled in fertility care programs for analysis. Only patients who gave written consent had their semen samples collected. Semen was collected by masturbation into a wide mouthed glass container after 3-4 days of sexual abstinence and was examined within 2 hours after each collection. After liquefaction, the semen was analyzed according to World Health Organization criteria [75].

During sample collection, the participant's age and information related to lifestyle (with attention to alcohol and tobacco consumption) were also collected. Clinical and laboratory studies were carried out in accordance with the recommendations of the manufacturers of diagnostic test kits and systems using modern laboratory technologies.

2.5. Data Analysis

Collected data were cleaned, entered, hardcoded and analyzed using Statistical Package for Social Sciences (SPSS) version 23. Summary statistics was presented using tables. Age was categorized and summarized with mean and standard deviation. Categorical variables were presented as proportions. Descriptive statistics were computed for relevant variables. The significant level of 5% (0.05) was set for all statistical procedures. Chi-square test (χ^2) was used to test the hypotheses and to compare the significance between lifestyle and its outcomes on semen quality at a level of significance set at $p < 0.05$.

2.6. Ethical Consideration

Ethical clearance was obtained from the Governing Board of Medlink Clinic. This ethical clearance ensured adequate professional work ethics and confidentiality of all patients' data. Signed consent from every participant was obtained. This research was also carried out with due observance of the ethical principles of the Declaration of Helsinki (DoH) in 2013 concerning human research.

3. Results

The findings in this study are presented in tables as follows: Table 1 shows the demographic distribution of the respondents. The mean age was 37.8 ± 10.3 years and the age bracket with the highest frequency was 28-37 years. With regards to their lifestyle, the participants were grouped into five. Frequent alcohol consumers, non-frequent alcohol consumers, non-alcohol consumers, tobacco smokers and non-smokers. Non-alcohol consumers and non-tobacco smokers accounted for 9.2% and 11.7% respectively. Alcohol consumers accounted for 38.8% of the respondents while tobacco smokers accounted for 40.3%.

Table 1. Demographic characteristics of participants, $n=196$.

Demographic characteristics	Options	Frequency, n (%)
Age range (years)	18-27	24 (12.2)
	28-37	80 (40.8)
	38-47	58 (29.6)
	48-57	21 (10.7)
	58-65	13 (6.6)
	Total	196 (100)
	Mean age = 37.8 ± 10.3	
Lifestyle	Frequent alcohol consumers	57 (29.1)
	Non-frequent alcohol consumers	19 (9.7)
	Non-alcohol consumers	18 (9.2)

Demographic characteristics	Options	Frequency, n (%)
	Tobacco smokers	79 (40.3)
	Non-smokers	23 (11.7)
	Total	196 (100)

Table 2 shows the distribution of the samples analyzed with only 18.4% of the participants having normospermia after sperm volume, count, motility, and morphological assessments. On sperm count, oligospermia (sperm count <20 million per ml of semen) was evident in 37.8% of the samples, cryptozoospermia (sperm count <100,000 per ml) was seen in 4.1% of the samples, while azoospermia (sperm count 0 per

ml i.e. absence of spermatozoa in semen) had the least frequency of 2.6%. On motility, 29.6% of the samples showed asthenozoospermia (<50% spermatozoa with forward progression i.e. poor sperm motility). On morphology, 7.7% showed teratozoospermia (<30% spermatozoa with normal morphology i.e. poor sperm morphology).

Table 2. Distribution of semen parameters of the samples analyzed based on sperm volume, count, motility, and morphology.

Semen parameters	Classification	Frequency, n (%)
Sperm volume, count, motility, morphology	Normospermia	36 (18.4)
Sperm count	Oligospermia	74 (37.8)
zz	Cryptozoospermia	8 (4.1)
	Azoospermia	5 (2.6)
Motility	Asthenozoospermia	58 (29.6)
Morphology	Teratozoospermia	15 (7.7)

Table 3 shows the distribution of semen qualities among the participants. Only twelve (21.1%) of the fifty-seven frequent alcohol consumers had normospermia, and asthenozoospermia accounted for the highest class of poor semen quality (77.84%). Ten (52.6%) of the non-frequent alcohol consumers were normospermic while 46.4% of them had poor semen quality. Surprisingly, 77.8% of the non-alcohol consumers had poor semen quality, while only 22.2% were normospermic. Most (89.9%) of the smokers had poor semen quality, with oligospermia accounting for the

highest occurrence (81.7%), and only 10.1% were normospermic. Sixteen (69.6%) of non-smokers had poor semen quality, while 30.4% had normospermia. Irrespective of the lifestyle, 20.9% of the participants were normospermic while 79.1% had poor semen quality. Out of the poor semen qualities, the most prevalent was oligospermia (47.7%), followed by asthenozoospermia (34.2%). The occurrence of cryptozoospermia and azoospermia was low at 5.2% and 3.2% respectively.

Table 3. Distribution of semen qualities among the study participants, n=196.

Participants' categories	Normo-spermia	Oligo-spermia	Cryptozoo-spermia	Azoo-spermia	Asthenozoospermia	Teratozoo-spermia	Total
Frequent alcohol consumers	12	5	1	1	35	3	57
Non-frequent alcohol consumers	10	0	1	1	5	2	19
Non-alcohol consumers	4	0	6	1	4	3	18
Smokers	8	58	0	2	6	5	79
Non-smokers	7	11	0	0	3	2	23
Total	41	74	8	5	53	15	196

3.1. Tests of Hypotheses

Three research hypotheses were tested in this study and all are displayed in Table 4. These hypotheses are to determine the relationship between lifestyle and semen quality at a level of significance $p < 0.05$.

3.2. Hypothesis One

H_0 : There is no significant relationship between alcohol/tobacco consumption and semen quality.

H_1 : There is a significant relationship between alcohol/tobacco consumption and semen quality.

The null hypothesis (H_0) was rejected because the p -value is less than 0.05 ($p = 0.024$), and the Chi-square test value (17.89) is greater than the critical value (5.99). This implies that there is a significant relationship between alcohol/tobacco

consumption and poor semen quality (male infertility).

3.3. Hypothesis Two

H_0 : There is no significant relationship between non-consumption of alcohol/tobacco and semen quality.

H_1 : There is a significant relationship between non-consumption of alcohol/tobacco and semen quality.

The null hypothesis (H_0) was accepted because the p -value is greater than 0.05 ($p = 0.56$), and the Chi-square test statistics value (0.35) is lesser than the critical value (3.84). This implies that there is no significant relationship between non-consumption of alcohol/tobacco and good semen quality (male fertility).

3.4. Hypothesis Three

H_0 : There is no significant relationship between alcohol

consumption and semen quality.

H₁: There is a significant relationship between alcohol consumption and semen quality.

The null hypothesis (H₀) was rejected because the *p*-value is

less than 0.05 (*p*=0.0086), and the Chi-square test statistics value (6.91) is greater than the critical value (3.84). This implies that there is a significant relationship between alcohol consumption and poor semen quality (male infertility).

Table 4. Statistical comparison between the occurrence of normospermia and poor sperm quality among the study participants based on their lifestyles.

Participants' categories	Normo-spermia	Poor sperm quality	Normo-spermia	Poor sperm quality	Total	<i>p</i> -value	<i>df</i>	critical value	χ^2
	Actual values		Expected values						
Alcohol and tobacco consumption									
Frequent alcohol consumers	12	45	11.03	45.97	57	0.024	2	5.99	17.89
Non-frequent alcohol consumers	10	9	3.68	15.32	19				
Smokers	8	71	15.29	63.71	79				
Total	30	125	30	125	155				
Non-alcohol and tobacco consumption						0.56	1	3.84	0.35
Non-alcohol consumers	4	14	4.83	13.17	18				
Non-smokers	7	16	6.17	16.83	23				
Total	11	30	11	30	41				
Alcohol consumption						0.0086	1	3.84	6.91
Frequent alcohol consumers	12	45	16.5	40.5	57				
Non-frequent alcohol consumers	10	9	5.5	13.5	19				
Total	22	54	22	54	76				

Abbreviation and symbol: *df* = degree of freedom; χ^2 = Chi-square test

4. Discussion

Infertility is becoming a topic of concern in the society [16], and it has led to the dissolution of several marriages in African communities. Males and females could be infertile and several studies have shown that lifestyle, pollutants, diet, and psychological stress could play a role in affecting semen parameters and causing male infertility [2, 6, 7, 16]. This study examined the lifestyle (alcohol and tobacco consumption) of Liberian men visiting a fertility clinic, and the relationship between these lifestyles and semen parameters. This therefore shows that the findings of this study do not interpret the overall rate of infertility among Liberian men.

The first discovery in this study is that 18.4% of the men who visited the fertility clinic for treatment were normospermic and 81.6% had poor sperm quality. These values are similar to the findings of Ali, *et al* that showed 13.2% normospermia and 86.8% abnormal semen parameters [3], but inversely related to the findings of Raj, *et al* where patients with normospermia accounted for 79.6% and oligospermic men were just 20.4% [52]. The incidence of the classes of poor semen quality varied among the categories of the participants. Of the parameters that cause poor semen quality, low sperm count was observed to be the most prevalent (54.4%), followed by sperm motility (36.2%) and lastly, sperm morphology (9.4%). However, the overall incidence of low sperm count among the men was 44.5%, a value close to the 44% in Abdullah, *et al* study on Sudanese patients [1], and higher than the 25% in Jajoo, *et al* study on Central Indian patients [42].

Low sperm count could be classified into oligospermia, cryptozoospermia, and azoospermia. In this study, oligospermia, cryptozoospermia, and azoospermia accounted for 37.8%, 4.1%, and 2.6% respectively. The incidence of

oligospermia in this study is close to the 32.1% reported by Ramya, *et al* [53], but higher than the 29.13% in Samal, *et al* [58]; 24.8% in Kalavathi, *et al* [44]; 20.4% in Raj, *et al* [52]; and 7.2% in Ali, *et al* [3] studies.

Alcohol and tobacco are globally abused and their adverse effects on health are deleterious. This study found that the category of men with the highest abnormality in semen parameters was tobacco smokers (89.9%) followed by frequent alcohol consumers (78.9%). This study also revealed that there is a relationship between the quantity of alcohol consumed and semen quality, a result similar to the findings of Amor, *et al* [4] and Raj, *et al* [52]. Raj, *et al* study analyzed the semen of 250 males attending a tertiary hospital in Maharashtra, India and found that only 8.3% frequent alcohol consumers and 36.4% non-frequent alcohol consumers were fertile. This present study found out that 21.1% frequent alcohol consumers and 52.6% non-frequent alcohol consumers were normospermic (fertile). This finding collaborates with the findings of Hansen, *et al* and Ricci, *et al* who reported that regular alcohol intake decreased semen volume and concentration of sperm [19, 54].

These effects of alcohol intake on semen quality have been reported to be caused by oxidative stress caused by an imbalance between Reactive Oxygen Species (ROS) produced by the alcohol consumed in the form of free radicals that contain one or more unpaired electrons and antioxidants [15, 62]. Frequent consumption of alcohol also stimulates lipid peroxidation that produces numerous electrophilic aldehydes, eg. malondialdehyde that can attack many cellular targets and cause DNA fragmentation [12, 15, 79, 81]. It also reduces the effects of antioxidant enzymes such as superoxide dismutase and glutathione [14].

With regards to the second category of participants of this study who are tobacco consumers, there exists a relationship between tobacco consumption and semen quality. This result

is also similar to the findings of Raj, *et al* where 0% frequent tobacco smokers, 41% non-frequent tobacco smokers, and 79.7% non-smokers were fertile [52]. This present study shows that 10.1% tobacco smokers and 30.4% non-smokers were normospermic (fertile). Amor, *et al*'s study presented similar findings with semen parameters significantly better in non-tobacco smokers than in tobacco smokers [4].

In addition, tobacco smoking has been documented to also cause oxidative stress due to the production of Reactive Oxygen Species (ROS) [46, 47], and its toxic effects are evident on sperm DNA [13], on sperm motility [57], and on sperm morphology [49].

The tests of hypotheses showed that there is a significant relationship ($p=0.024$) between alcohol and tobacco consumption and poor semen quality (male infertility). There is no significant relationship ($p=0.56$) between non-consumption of alcohol/tobacco and good semen quality (male fertility). There is also a significant relationship ($p=0.0086$) between alcohol consumption only and poor semen quality (male infertility).

5. Conclusion

Alcohol and tobacco consumption reduce male sperm quality, but abstinence from both does not guarantee optimum semen quality. Alcohol and/or tobacco consumption adversely affect the semen parameters (sperm volume, count, motility, and morphology). Alcohol and tobacco consumption combined with bad lifestyle habits affect semen quality. Therefore, it is safer to abstain from alcohol and/or tobacco consumption since it has been proven that the adverse effects on sperm quality are directly proportional to the consumption rate of either or both substances. It is recommended that health institutions and healthcare workers should provide rigorous health education and awareness campaigns to the male populace on the negative impacts of alcohol and tobacco consumption aimed at not only improving male fertility, but also to improve life expectancy especially among the male population. The direction of future research on semen analysis should be to study other sperm abnormalities such as aspermia (no ejaculate), and asthenoteratozoospermia (combination of abnormal sperm motility, morphology and count).

Conflict of Interest

The authors guarantee responsibility for everything published in this manuscript, as well as the absence of a conflict of interest and the absence of their financial interest in performing this research and writing this manuscript. This manuscript was written from an original research work and has never been published, neither is it under consideration for publication elsewhere.

Acknowledgements

Special thanks to the staff of Medlink Clinic, Monrovia,

Liberia; and to all the men who participated in this study for their cooperation and support in carrying out this research.

References

- [1] Abdullah AA, Ahmed M, Oladokun A. Prevalence of infertility in Sudan: A systematic review and meta-analysis. *Qatar Medical Journal*. 2021; 2021 (3): 47-57.
- [2] Afeiche M, Williams PL, Mendiola J, Gaskins AJ, Jørgensen N, Swan SH, *et al*. Dairy food intake in relation to semen quality and reproductive hormone levels among physically active young men. *Human Reproduction*. 2013; 28 (8): 2265-2275.
- [3] Ali AFM, Modawe G, Rida MA, Abdrabo AA. Prevalence of abnormal semen parameters among male patients attending the fertility center in Khartoum, Sudan. *Journal of Medical and Life Science*. 2022; 4 (1): 1-8.
- [4] Amor H, Hammadeh ME, Mohd I, Jankowski PM. Impact of heavy alcohol consumption and cigarette smoking on sperm DNA integrity. *Andrologia*. 2022; 54 (7): e14434.
- [5] Amor H, Zeyad A, Hammadeh ME. Tobacco smoking and its impact on the expression level of sperm nuclear protein genes: H2BFWT, TNP1, TNP2, PRM1 and PRM2. *Andrologia*. 2021; 53: e13964. DOI: 10.1111/and.13964.
- [6] Barazani Y, Katz BF, Nagler HM, Stember DS. Lifestyle, environment, and male reproductive health. *Urologic Clinics of North America*. 2014; 41 (1): 55-66.
- [7] Boeri L, Capogrosso P, Ventimiglia E, Pederzoli F, Cazzaniga W, Chierigo F, *et al*. Heavy cigarette smoking and alcohol consumption are associated with impaired sperm parameters in primary infertile men. *Asian Journal of Andrology*. 2019; 21 (5): 478-485. DOI: 10.4103/aja.aja_110_18.
- [8] Center for Disease Control and Prevention. General alcohol information. *Fact Sheet*. Atlanta, GA: CDC, 2005. Available online: www.cdc.gov/alcohol/factsheets/general_information.htm
- [9] Center for Disease Control and Prevention. The health consequences of smoking: A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2004.
- [10] Clemens KJ, Caillé S, Stinus L, Cador M. The addition of five minor tobacco alkaloids increases nicotine-induced hyperactivity, sensitization and intravenous self-administration in rats. *International Journal of Neuropsychopharmacology*. 2009; 12 (10): 1355-1366.
- [11] Condorelli RA, Calogero AE, Vicari E, La Vignera S. Chronic consumption of alcohol and sperm parameters: our experience and the main evidences. *Andrologia*. 2015; 47 (4): 368-379. DOI: 10.1111/and.12284.
- [12] Conklin KA. Dietary antioxidants during cancer chemotherapy: impact on chemotherapeutic effectiveness and development of side effects. *Nutrition and Cancer*. 2000; 37 (1): 1-18. DOI: 10.1207/S15327914NC3701_1.
- [13] Cui X, Jing X, Wu X, Wang Z, Li Q. Potential effect of smoking on semen quality through DNA damage and the downregulation of Chk1 in sperm. *Molecular Medicine Reports*. 2016; 14 (1): 753-761. DOI: 10.3892/MMR.2016.5318/HTML.

- [14] Doshi SB, Khullar K, Sharma RK, Agarwal A. Role of reactive nitrogen species in male infertility. *Reproductive Biology and Endocrinology*. 2012; 10: 1–11.
- [15] Ekechi HO, Ikwuka AO, Udeh FC, Abraham JC. Effects of ethanol extract of *Rauwolfia vomitoria* leaf on lipid profile and cerebellar histology in cisplatin-induced oxidative stress. *British Journal of Medical and Health Research*. 2023; 10 (5): 16-39. DOI: 10.5281/zenodo.8042521.
- [16] Fang YY, Wu QJ, Zhang TN, Wang TR, Shen ZQ, Jiao J, *et al.* Assessment of the development of assisted reproductive technology in Liaoning province of China, from 2012 to 2016. *BMC Health Services Research*. 2018; 18 (1): 873. DOI: 10.1186/s12913-018-35.
- [17] Halmenschlager G, Rossetto S, Lara GM, Rhoden EL. Evaluation of the effects of cigarette smoking on testosterone levels in adult men. *Journal of Sexual Medicine*. 2009; 6 (6): 1763–1772. DOI: 10.1111/j.1743-6109.2009.01227.x.
- [18] Hamad MF, Shelko N, Kartarius S, Montenarh M, Hammadeh ME. Impact of cigarette smoking on histone (H2B) to protamine ratio in human spermatozoa and its relation to sperm parameters. *Andrology*. 2014; 2: 666–677. DOI: 10.1111/j.2047-2927.2014.00245.x.
- [19] Hansen ML, Thulstrup AM, Bonde JP, Olsen J, Hakonsen LB, Ramlau-Hansen CH. Does last week's alcohol intake affect semen quality or reproductive hormones? A cross-sectional study among healthy young Danish men. *Reproductive Toxicology*. 2012; 34: 457–462.
- [20] Heishman SJ, Kozlowski LT, Henningfield JE. Nicotine addiction: Implication for public health policy. *Journal of Social Issues*. 2010; 53 (1): 13-33.
- [21] Ikwuka AO. Risk factors for the pathogenesis of diabetes mellitus type 2. *Materials of 84th Scientific and Practical Conference of Students and Young Scientists with International Participation "Innovations in medicine"*. 2015; p. 19. Available online: http://www.ifnmu.edu.ua/images/snt/files/konferenciya/Tezu_2015.pdf
- [22] Ikwuka AO. Dyslipidemia risk severity in patients with diabetes mellitus type 2 and essential hypertension. *Journal of the 21st International Medical Congress of Students and Young Scientists*. 2017; p. 59.
- [23] Ikwuka AO. Effectiveness of dapagliflozin in patients with diabetes mellitus type 2 and essential hypertension. *Book of abstracts of the 7th International Students' Scientific Conference of Young Medical Researchers*. 2017; p. 102. Available online: http://www.stn.umed.wroc.pl/files/lm/Accepted_papers.16113.pdf
- [24] Ikwuka AO. Influence of dyslipidemia in patients with diabetes mellitus type 2 and essential hypertension. *The Pharma Innovation Journal*. 2017; 6 (3): 101-103. Available online: <http://www.thepharmajournal.com/archives/?year=2017&vol=6&issue=3&part=B>
- [25] Ikwuka AO, Haman IO. Features of kidney damage in patients with diabetes mellitus type 2 and essential hypertension. *Journal of 86th Scientific and Practical Conference of Students and Young Scientists with International Participation "Innovations in medicine"*. 2017; p. 144. Available online: [http://www.ifnmu.edu.ua/images/snt/86-konf-tezi%20\(1\).pdf](http://www.ifnmu.edu.ua/images/snt/86-konf-tezi%20(1).pdf)
- [26] Ikwuka AO, Virstyuk NG, Luchko OR. Features of the functional state of kidneys in patients with diabetes mellitus type 2 and essential arterial hypertension. *Materials of scientific-practical conference with international participation "Babinkivski reading"*. 2017; p. 48.
- [27] Ikwuka AO. Clinical dynamics in patients with diabetes mellitus type 2 and concomitant essential hypertensive disease treated with dapagliflozin. *Journal of the 22nd International Medical Congress of Students and Young Scientists*. 2018; p. 32.
- [28] Ikwuka AO. Clinical effectiveness of SGLT-2 inhibitors in patients with diabetes mellitus type 2 and essential hypertensive disease. *Endocrine Practice*. 2018; 24 (1): 74. DOI: 10.1016/S1530-891X(20)47129-0.
- [29] Ikwuka AO. Features of kidney damage in patients with arterial hypertension and type 2 diabetes mellitus and optimization of treatment. *Specialized Academic Council IFN MU*. 2018; Available online: http://www.ifnmu.edu.ua/images/zagalna_informacia/spec_vcheni_radi/2017-2019/%D0%9420.601.01/Ikwuka/Avtoreferat.pdf
- [30] Ikwuka AO, Paliy Yu. Structural changes of the left ventricular myocardium in patients with essential arterial hypertension and diabetes mellitus type 2. *Abstracts of the 87th Scientific Conference of Students and Young Scientists with International Participation "Innovations in medicine"*. 2018; p. 25-26. Available online: https://www.ifnmu.edu.ua/images/snt/zaproshennia_eng.pdf
- [31] Ikwuka AO. Clinical dynamics of nephropathy in patients with diabetes mellitus type 2 and concomitant essential hypertensive disease. *Clinical Medicine*. 2019; 19 (2): s39. DOI: 10.7861/clinmedicine.19-2-s39.
- [32] Ikwuka AO. Clinical effectiveness of GLP-1 RAs in patients with metabolic syndrome diseases. *Endocrine Practice*. 2019; 25 (1): 104-105. DOI: 10.1016/S1530-891X(20)46611-X.
- [33] Ikwuka AO. *Dr. Aloy's Core Essential Series (DACES) Immunology*. 1st Edition. Science and Education Publishing, USA. 2023; p. 31. ISBN: 978-978-795-866-7.
- [34] Ikwuka AO. *Dr. Aloy's Core Essential Series (DACES) Medical Genetics*. 1st Edition. Science and Education Publishing, USA. 2023; p. 9. ISBN: 978-1-958293-02-7.
- [35] Ikwuka AO, Virstyuk NG. Pattern of cardiac remodelling of the left ventricle in patients with essential hypertensive disease and concomitant type 2 diabetes mellitus. *Clinical Medicine*. 2019; 19 (3): s92. DOI: 10.7861/clinmedicine.19-3-s92.
- [36] Ikwuka AO, Virstyuk NG. Influence of SGLT2 inhibitor and A2RB (AT1) on fibrogenesis and heart failure in patients with essential hypertensive disease combined with diabetes mellitus type 2. *E-Poster No. 143 of the 44th & 45th Annual General and Scientific Meeting of the West African College of Physicians (WACP), 1 - 3 November, 2021*. 2021; DOI: 10.13140/RG.2.2.26912.87047.
- [37] Ikwuka AO, Virstyuk N. Prognostic markers of nephropathy in patients with dual metabolic syndrome diseases (essential hypertensive disease and concomitant type 2 diabetes mellitus). *Endocrine Practice*. 2022; 28 (5): S65-S66. DOI: 10.1016/j.eprac.2022.03.164.
- [38] Ikwuka AO, Virstyuk N. Patterns and Influence of Cardio-Metabolic Insufficiency in Patients with Essential Hypertensive Disease and Concomitant Type 2 Diabetes Mellitus. *Endocrine Practice*. 2023; 29 (5): S32-S33. DOI: 10.1016/j.eprac.2023.03.076.

- [39] Ikwuka AO, Virstyuk NG, Luchko OR, Kobitovych I. Heterogeneity Of Renal Pathogenicity On The Background Of Asymptomatic Hyperuricemia In Patients With Dual Metabolic Syndrome Diseases (Essential Hypertensive Disease and Type 2 Diabetes Mellitus). *British Journal of Medical and Health Research*. 2023; 10 (2): 1-9. DOI: 10.5281/zenodo.7690636.
- [40] International Agency for Research on Cancer Working Group. Smokeless tobacco and some tobacco-specific N-nitrosamines. *World Health Organization Monographs on the Evaluation of Carcinogenic Risks to Humans*. 2007; 89: 345-346.
- [41] Irvine DS. Epidemiology and aetiology of male infertility. *Human Reproduction*. 1998; 13 (1): 33-44.
- [42] Jajoo S, Kalyani KR. Prevalence of abnormal semen analysis in patients of infertility at a rural set up in central India. *International Journal of Reproduction, Contraception, Obstetrics, and Gynecology*. 2013; 2: 161-164.
- [43] Jorsaraei SGA, Shibahara H, Ayustawati HY, Shiraishi Y, Khalatbari A, Pasha YY, *et al*. The in-vitro effects of nicotine, cotinine and leptin on sperm parameters analyzed by CASA system. *Iranian Journal of Reproductive Medicine*. 2008; 6 (3): 157-165.
- [44] Kalavathi DB. Male factor in infertility: Study from a tertiary care hospital. *International Journal of Reproduction, Contraception, Obstetrics, and Gynecology*. 2016; 5 (6): 2022-2025.
- [45] Kamal K. Monograph of tobacco (*Nicotianatabacum*). *Indian Journal of Drugs*. 2014; 2 (1): 5-23.
- [46] Kumar SB, Chawla B, Bisht S, Yadav RK, Dada R. Tobacco use increases oxidative DNA damage in sperm - possible etiology of childhood cancer. *Asian Pacific Journal of Cancer Prevention*. 2015; 16: 6967-6972. DOI: 10.7314/APJCP.2015.16.16.6967.
- [47] La Maestra S, De Flora S, Micale RT. Effect of cigarette smoke on DNA damage, oxidative stress, and morphological alterations in mouse testis and spermatozoa. *International Journal of Hygiene and Environmental Health*. 2015; 218: 117-122.
- [48] Majo J, Ghezzi H, Cosio MG. Lymphocyte population and apoptosis in the lungs of smokers and their relation to emphysema. *European Respiratory Journal*. 2001; 17 (5): 946-953. DOI: 10.1183/09031936.01.17509460.
- [49] Mostafa T. Cigarette smoking and male infertility. *Journal of Advanced Research*. 2010; 1 (3): 179-186. DOI: 10.1016/J.JARE.2010.05.002.
- [50] Muthusami KR, Chinnaswamy P. Effect of chronic alcoholism on male fertility hormones and semen quality. *Fertility and Sterility*. 2005; 84: 919-924. DOI: 10.1016/j.fertnstert.2005.04.025.
- [51] Pant N, Kumar G, Upadhyay AD, Patel DK, Gupta YK, Chaturvedi PK. Reproductive toxicity of lead, cadmium, and phthalate exposure in men. *Environmental Science and Pollution Research*. 2014; 21 (18): 11066-11074. DOI: 10.1007/s11356-014-2986-5.
- [52] Raj A, Iqbal B, Sharma A, Gore CR, Kumar H, Singh M. Effect of alcohol and smoking on semen analysis parameters. *Medical Journal of Dr. D. Y. Patil Vidyapeeth*. 2022; 15 (5): 696-700.
- [53] Ramya C, Renuka IV, Premalatha P, Madhavi K. Patterns of semen analysis in male partners of infertile couples at a tertiary care hospital. *Indian Journal of Pathology and Oncology*. 2017; 4 (4): 536-539.
- [54] Ricci E, Noli S, Ferrari S, La Vecchia I, Cipriani S, De Cosmi V, *et al*. Alcohol intake and semen variables: cross-sectional analysis of a prospective cohort study of men referring to an Italian Fertility Clinic. *Andrology*. 2018; 6 (5): 690-696.
- [55] Sadler TW. *Langman's Medical Embryology*. 12th Edition. Lippincott Williams & Wilkin. 2012; p. 36-42.
- [56] Saha S, Roy P, Corbitt C, Kakar SS. Application of stem cell therapy for infertility. *Cells*. 2021; 10: 1613. DOI: 10.3390/cells10071613.
- [57] Saleh RA, Agarwal A, Sharma RK, Said TM, Sikka SC, Thomas AJ. Evaluation of nuclear DNA damage in spermatozoa from infertile men with varicocele. *Fertility and Sterility*. 2003; 80 (6): 1431-1436. DOI: 10.1016/S0015-0282(03)02211-8.
- [58] Samal S, Dhadwe K, Gupta U, Gupta NK. Epidemiological study of male infertility. *Indian Medical Gazette*. 2012; 6 (5): 174-180.
- [59] Sengupta P, Dutta S, Krajewska-Kulak E. The disappearing sperms: Analysis of reports published between 1980 and 2015. *American Journal of Men's Health*. 2017; 11 (4): 1279-1304. DOI: 10.1177/1557988316643383.
- [60] The World Bank. Data: Prevalence of current tobacco use, males (% of male adults)-Liberia. 2023. Available online: <https://data.worldbank.org/indicator/SH.PR.V.SMOK.MA?locations=LR>
- [61] The World Bank. Data: Total alcohol consumption per capita (liters of pure alcohol, projected estimates, 15+ years of age) - Liberia. 2023. Available online: <https://data.worldbank.org/indicator/SH.ALC.PCAP.LI?locations=LR>
- [62] Tsermpini EE, Plemenitaš Ilješ A, Dolžan V. Alcohol-induced oxidative stress and the role of antioxidants in alcohol use disorder: A systematic review. *Antioxidants*. 2022; 11: 1374-1407. DOI: 10.3390/antiox11071374.
- [63] Udeh FC, Ikwuka AO, Epete MA, Igwe EC. Effects of local tobacco snuff ingestion during pregnancy on renal functions and histology architecture of female Wistar rats and on the birth weight of their pups. *American Journal of Medical Sciences and Medicine*. 2023; 11 (1): 1-5. DOI: 10.12691/ajmsm-11-1-1.
- [64] Udeh FC, Ikwuka AO, Epete MA, Igwe EC. Effects of oral consumption of *Nicotianatabacum* during pregnancy on the liver and prolactin levels of adult female Wistar rats. *European Journal of Veterinary Medicine*. 2023; 3 (2): 1-5. DOI: 10.24018/ejvetmed.2023.3.2.93.
- [65] Virstyuk NG, Ikwuka AO, Haman IO, Adebomi MS. Diabetes mellitus type 2, arterial hypertension and dyslipidemia. *Materials of 2nd International Scientific and Practical Conference "Therapeutic readings: modern aspects of diagnosis and treatment of diseases of internal organs"*. 2016; p. 46-47.
- [66] Virstyuk NG, Ikwuka AO. Diagnostic and prognostic markers of the diabetes mellitus type 2 course in connection with essential arterial hypertension taking into account the kidney function. *Precarpathian Journal Pulse* (ISSN: 2304-7437). 2017; 8 (44): 53-62.

- [67] Virstyuk NG, Ikwuka AO. Features of asymptomatic hyperuricemia in patients with diabetes mellitus type 2 and concomitant essential arterial hypertension. *Clinical and Experimental Pathology*. 2018; 1 (63): 22-26. DOI: 10.24061/1727-4338.XVII.1.63.2018.5.
- [68] Virstyuk NG, Ikwuka AO. Nephropathic characteristics in patients with diabetes mellitus type 2 and essential hypertensive disease. *Art of Medicine*. 2019; 1 (5): 44-47. DOI: 10.21802/artm.2019.1.9.44.
- [69] Virstyuk NG, Ikwuka AO. Asymptomatic hyperuricemia and functional state of the kidneys in patients with essential arterial hypertension and concomitant diabetes mellitus type 2. *European Journal of Clinical Medicine*. 2021; 2 (3): 100-104. DOI: 10.24018/clinicmed.2021.2.3.65.
- [70] Virstyuk NG, Ikwuka AO, Didushko OM. Effect of dapagliflozin on the level of uric acid during asymptomatic hyperuricemia in patients with diabetes mellitus type 2 and concomitant arterial hypertension. *Art of Medicine*. 2018; 1 (5): 21-26. Available online: <https://art-of-medicine.ifnmu.edu.ua/index.php/aom/article/view/179/150>
- [71] Virstyuk NH, Ikwuka AO. Dapagliflozin influence on the clinical course of diabetes mellitus type 2 and essential hypertension in patients. *Recent Advances in Environmental Science from the Euro-Mediterranean and Surrounding Regions*. Springer International Publishing AG. 2018; p. 2007-2008. DOI: 10.1007/978-3-319-70548-4_582.
- [72] Virstyuk NH, Ikwuka AO, Losyuk LV, Kobrynska OYa, Markiv HD. Dapagliflozin utility in patients with diabetes mellitus type 2 and essential hypertensive disease. *Actual Problems of Modern Medicine*. 2017; 4 (60) 1: 76-79. Available online: http://www.umsa.edu.ua/journal2stat4_2017_eng.html
- [73] Virstyuk NH, Ikwuka AO, Luchko OR, Kocherzhat OI. Peculiarities of renal insufficiency in patients with diabetes mellitus type 2 and arterial hypertension. *Materials of scientific-practical conference with international participation "Achievements and prospects of experimental and clinical endocrinology" Twentieth Danilevsky readings*. 2021; p. 86-87.
- [74] World Health Organization. *Fertility*. 2020; Available online: <https://www.who.int/news-room/fact-sheets/detail/infertility>
- [75] World Health Organization. Standard Procedures. In: Cooper TG, editor. *WHO Laboratory Manual for the Examination and Processing of Human Semen*. 5th ed. Geneva. World Health Organization. 2010; p. 1-141.
- [76] World Health Organization. *The World Health Observatory*. 2023; Available online: <https://www.who.int/data/gho/data/themes/global-information-system-on-alcohol-and-health>
- [77] World Health Organization. Tobacco. *Fact Sheets*. 2022; Available online: <https://www.who.int/news-room/fact-sheets/detail/tobacco>
- [78] Wright J, Duchesne C, Sabourin S, Bissonnette F, Benoit J, Girard Y. Psychosocial distress and infertility: men and women respond differently *in vitro*. *Fertility and Sterility*. 1991; 55: 100-108.
- [79] Wu D, Cederbaum AI. Alcohol, oxidative stress, and free radical damage. *Alcohol Research and Health*. 2003; 27 (4): 277–284. DOI: 10.1079/pns2006496.
- [80] Zegers-Hochschild F, Adamson GD, de Mouzon J, Ishihara O, Mansour R, Nygren K, *et al.* The International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary on ART terminology, 2009. *Human Reproduction*. 2009; 24: 2683-2687.
- [81] Zorn B, Vidmar G, Meden-Vrtovec H. Seminal reactive oxygen species as predictors of fertilization, embryo quality and pregnancy rates after conventional *in vitro* fertilization and intracytoplasmic sperm injection. *International Journal of Andrology*. 2003; 26 (5): 279–285. DOI: 10.1046/j.1365-2605.2003.00424.x.