

Review Article

Joint Hypermobility Syndrome: A Narrative Review

Jobson Lopes de Oliveira^{1,2,*}, Igor Albuquerque Nogueira¹, Camila Távora Nogueira², Saulo Venicius Parente Lopes³, Gabriela Silva Teles⁴, Raul Sancho de Carvalho Rocha², Pedro Iughetti Morais², Ana Beatriz Cordeiro Carvalho², Pedro Henrique de Santana Pereira², David Johnson Pinheiro Alencar²

¹Department of Internal Medicine, Faculty of Medicine, Federal University of Ceará, Fortaleza, Brazil

²Department of Internal Medicine, Faculty of Medicine, Christus University Center (Unichristus), Fortaleza, Brazil

³Department of Internal Medicine, Faculty of Medicine, University of Fortaleza (Unifor), Fortaleza, Brazil

⁴Albert Sabin Children's Hospital, Fortaleza, Brazil

Email address:

jobsonlopesdeoliveira@gmail.com (Jobson Lopes de Oliveira)

*Corresponding author

To cite this article:

Jobson Lopes de Oliveira, Igor Albuquerque Nogueira, Camila Távora Nogueira, Saulo Venicius Parente Lopes, Gabriela Silva Teles, Raul Sancho de Carvalho Rocha, Pedro Iughetti Morais, Ana Beatriz Cordeiro Carvalho, Pedro Henrique de Santana Pereira, David Johnson Pinheiro Alencar. Joint Hypermobility Syndrome: A Narrative Review. *American Journal of Internal Medicine*. Vol. 11, No. 3, 2023, pp. 35-40. doi: 10.11648/j.ajim.20231103.12

Received: May 1, 2023; **Accepted:** May 15, 2023; **Published:** May 25, 2023

Abstract: Joint hypermobility syndrome (JHS) is a complex connective tissue disorder characterized by excessive joint mobility and chronic musculoskeletal pain. The condition extends beyond the joints and can affect various systems such as the skin, gastrointestinal tract, neurological system, and cardiovascular system, necessitating a multidisciplinary approach to optimize care. Notably, the prevalence of JHS is higher in young individuals, females, and those of Asian descent. Despite being a common condition, JHS is frequently underdiagnosed, resulting in persistent pain and disability. In order to diagnose JHS, it is necessary to rule out the presence of any consistent feature that indicates the existence of other connective tissue disorders that partially overlap with it, including Marfan and Ehlers-Danlos syndromes. A thorough clinical assessment, coupled with the application of the Brighton Criteria, can aid in making an accurate diagnosis. Management of JHS poses significant challenges and typically involves symptomatic treatment, including physiotherapy, rehabilitation, and pharmacological therapy. Additionally, a multidisciplinary approach, including collaboration with different specialists, is vital. Further research is needed to improve our understanding of JHS and develop interventions based on solid evidence. However, with comprehensive management strategies, individuals with JHS can achieve better pain control and improved function, enabling them to lead more fulfilling lives.

Keywords: Hypermobility Syndrome, Ehlers-Danlos Syndrome, Joint Pain, Chronic Pain

1. Introduction

Joint hypermobility syndrome (JHS) is a heritable connective tissue disorder characterized by hyperextensibility of joints and chronic musculoskeletal pain. This condition affects the articulations of all four limbs and the axial skeleton [1]. However, systemic complaints, such as dysautonomia, fatigue, affective disorders, and gastrointestinal manifestations, are also commonly reported [2, 3]. Despite

having a prevalence similar to that of other rheumatologic diseases such as fibromyalgia, gout, and rheumatoid arthritis, JHS is often underdiagnosed [4, 5]. In this narrative review, we will discuss the diverse manifestations of JHS, along with epidemiological aspects, pathophysiological mechanisms, diagnostic approaches, and treatment practices.

2. Epidemiology

Generalized joint hypermobility refers to an extraordinary flexibility in joint movements that affects both large and small peripheral joints, as well as the spine [6]. This condition has a significant genetic component, with a heritability estimate of 70% in females based on a twin study [7]. It is a multifactorial trait that can affect individuals of all genders, with prevalence rates ranging between 6% to 57% in women and 2% to 35% in men, and these rates can vary depending on age and ethnicity [8, 9].

JHS is more common in women. Among individuals of different ethnic origins, the highest prevalence of the condition is seen in individuals of Asian descent, followed by those of African and European descent, respectively. It can affect individuals of all age groups, with milder forms typically occurring in older patients, while more severe cases are often seen in younger individuals. In pediatric patients, JHS is present in 10 to 15% of children with chronic pain syndrome [1, 2, 5].

3. Pathophysiology

The pathophysiological mechanisms of JHS involve both genetic and environmental factors. Although several genes have been linked to the condition, their penetrance varies, and the influence of individual genes on the phenotypes is not well understood due to polygenic heterogeneity. Relevant polymorphisms have been identified in genes such as *TNXB*, *COL3A1*, and *LZTS1*, but further research is needed to clarify the role of genetic diversity in JHS [3].

Environmental factors such as weight gain, trauma, and surgical procedures also contribute to the development of unspecific articular hypermobility. These factors can lead to structural changes in joints, muscles, and connective tissue, resulting in chronic pain and altered proprioception [9]. The formation of micro-lesions in connective tissue may also facilitate joint fragility and structural damage, leading to a chain of events that culminates in chronic and debilitating pain [10].

The interaction between genetic predisposition and environmental factors gives rise to a wide range of JHS manifestations, which extend beyond musculoskeletal symptoms. A better understanding of the mechanisms underlying JHS is necessary to improve clinical management and potentially prevent the condition [11].

4. Diagnosis

Diagnosing JHS involves a thorough clinical evaluation. There are various connective tissue disorders that show general joint hypermobility, such as Marfan or Ehlers-Danlos syndromes (EDS). Since JHS is a diagnosis made by excluding other potential diseases, it is crucial to identify any other symptoms and carefully inquire about the family's medical history regarding connective tissue disorders. The symptoms of JHS and EDS-hypermobility type (EDS-HT) or type 3 are very similar. While other symptoms may aid in diagnosing non-EDS disorders, generalized joint hypermobility is usually the primary indicator of EDS. Table 1 provides a list of symptoms that can help differentiate between different disorders. [3, 5, 12, 13].

Table 1. Differential diagnosis of joint hypermobility syndrome.

DISEASE	CLINICAL FEATURES
Ehlers-Danlos syndrome (mainly type 3)	Stretchy, thin, soft, and velvety skin Positive family history Unexplained striae Generalized joint hypermobility Recurrent dislocations High stature
Marfan syndrome	Arachnodactyly Ectopia lens Aortic root dilation and aortic aneurysms Pectus excavatum Low stature
Osteogenesis imperfecta	Bluish sclerae Multiple fractures Sensorineural hearing loss Dental manifestations
Loeys-Dietz syndrome	Aortic aneurysms with dissection risk Hypertelorism Bifid uvula / Cleft palate

To standardize the assessment of generalized joint hypermobility, Beighton's score (Figure 1) was developed [14]. It consists of five simple maneuvers that measure the degree of hypermobility of each joint using a nine-point scoring system. A score of four or greater is considered indicative of generalized joint hypermobility [15], which is one of the major criteria for JHS according to Brighton's

criteria (Table 2). The other major criterion is the presence of arthralgia in more than four joints that lasts longer than three months [16].

Additionally, a five-part questionnaire (Table 2) has been developed to screen for JHS [17]. A positive response to at least two of the five questions suggests a greater likelihood of joint hypermobility syndrome and warrants further

investigation. This tool is especially useful for primary care physicians due to its simplicity and objectivity [10, 17].






MANEUVER	EXAMPLE	NUMBER OF POINTS
Hyperextension of the elbows to more than 10 degrees		Each side made equals 1 point (maximum 2 points)
Hyperextension of the fifth finger		Each side made equals 1 point (maximum 2 points)
Hyperextension of the trunk with the palms of the hands touching the ground and with the knees extended		1 point if done
Knee hyperextension to more than 10 degrees		Each side made equals 1 point (maximum 2 points)
Juxtaposition of thumb and forearm		Each side made equals 1 point (maximum 2 points)

Figure 1. The Beighton scoring system. Each of the above maneuvers earns 1 point, with a maximum possible score of 9 points. Scoring 4 or more points indicates the presence of generalized joint hypermobility.

Table 2. Brighton criteria for assessment of JHS and 5-Point Questionnaire for JHM.

Brighton criteria for assessment of JHS		5-Point Questionnaire for JHM
MAJOR CRITERIA	CONSIDERATIONS	
Beighton score $\geq 4/9$	Refer to Figure 1 for the calculation of the score.	1) Can you now (or could you ever) place your hands flat on the floor without bending your knees?
Arthralgias	For longer than 3 months, in more than 4 joints.	

important to note that it is not established whether generalized exercises or specific group training work best for the condition [32, 33]. Regular exercise has been shown to provide relief from JHS symptoms. However, excessive joint movement, strenuous training, and excessive focus on articular flexibility may worsen symptoms and lead to injuries. Therefore, exercises that focus on balance and gentle stretching of joints may be more appropriate [34, 35].

Pharmacological measures such as oral analgesics, selective serotonin reuptake inhibitors, anticonvulsants, and tricyclic antidepressants may be helpful, but caution and short-term therapy are advised [36]. Primary healthcare physicians should be educated about the clinical presentation of the condition to facilitate timely diagnosis and management. Referral to a specialist is indicated in case of systemic symptoms, such as urinary dysfunction, gastrointestinal manifestations, and dysautonomia [10].

7. Conclusions

JHS is a prevalent condition that often presents as chronic pain and must be distinguished from other collagenopathies. Unfortunately, it is an underdiagnosed condition that carries a stigma due to a lack of awareness regarding its symptoms and prognosis. JHS can affect multiple systems beyond joints. Thus, physicians' awareness plays a crucial role in early disease detection, and continuing medical education should be promoted. A multidisciplinary approach that encompasses the integrality of care and patient education is essential when dealing with JHS. Therapeutics must be individualized and based on longitudinal monitoring, taking into account each patient's priorities. Lifestyle changes, physiotherapy, and physical exercise are the most important elements of care, while psychotherapy and self-care techniques are also supported. However, there is a lack of evidence regarding risk factors, prevalence, and specific interventions, indicating the need for further studies to assess the true impact of JHS and develop better practices.

Abbreviations

JHS: joint hypermobility syndrome
EDS: Ehlers-Danlos syndromes
GI: gastrointestinal
MVP: mitral valve prolapse

Author's Contributions

JLO conceived the article. Data acquisition, analysis and/or interpretation: SVPL, GST, RSCR, PIM, ABCC, PHSP, DJPA. Draft and revision of the work: IAN, CTN, JLO. All authors approved the final version of the manuscript.

Consent for Publication

Written informed consent was obtained from the patient to

publish the image in this article. A copy of the written consent is available for review by the editor of this journal.

Competing Interests

The author declares that there are no competing interests.

References

- [1] Fikree A, Aziz Q, Grahame R. Joint hypermobility syndrome. *Rheum Dis Clin North Am*. 2013; 39 (2): 419-430.
- [2] Carbonell-Bobadilla N, Rodríguez-Álvarez A, Rojas-García G, Barragán-Garfias J, Orrantia-Vertiz M, Rodríguez-Romo R. Joint hypermobility syndrome. *Acta Ortopedica Mexicana*. 2020; 34 (6): 441-9.
- [3] Tinkle BT, Levy HP. Symptomatic Joint Hypermobility: The Hypermobile Type of Ehlers-Danlos Syndrome and the Hypermobility Spectrum Disorders. *Med Clin North Am*. 2019; 103 (6): 1021-1033.
- [4] Grahame R. Hypermobility: an important but often neglected area within rheumatology. *Nat Clin Pract Rheumatol*. 2008; 4 (10): 522-524.
- [5] Kumar B, Lenert P. Joint Hypermobility Syndrome: Recognizing a Commonly Overlooked Cause of Chronic Pain. *Am J Med*. 2017; 130 (6): 640-647.
- [6] Tinkle BT. Symptomatic joint hypermobility. *Best Pract Res Clin Rheumatol*. 2020; 34 (3): 101508.
- [7] Hakim AJ, Cherkas LF, Grahame R, Spector TD, MacGregor AJ. The genetic epidemiology of joint hypermobility: a population study of female twins. *Arthritis Rheum*. 2004; 50 (8): 2640-2644.
- [8] Remvig L, Jensen DV, Ward RC. Epidemiology of general joint hypermobility and basis for the proposed criteria for benign joint hypermobility syndrome: review of the literature. *J Rheumatol*. 2007; 34: 804-9.
- [9] Syx D, De Wandele I, Rombaut L, Malfait F. Hypermobility, the Ehlers-Danlos syndromes and chronic pain. *Clin Exp Rheumatol*. 2017; 35 Suppl 107 (5): 116-122.
- [10] Atwell K, Michael W, Dubey J, et al. Diagnosis and Management of Hypermobility Spectrum Disorders in Primary Care. *J Am Board Fam Med*. 2021; 34 (4): 838-848.
- [11] Tinkle B, Castori M, Berglund B, et al. Hypermobile Ehlers-Danlos syndrome (a.k.a. Ehlers-Danlos syndrome Type III and Ehlers-Danlos syndrome hypermobility type): Clinical description and natural history. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 48-69.
- [12] Pacey V, Tofts L, Wesley A, Collins F, Singh-Grewal D. Joint hypermobility syndrome: a review for clinicians. *J Paediatr Child Health*. 2015; 51 (4): 373-380.
- [13] Colombi M, Dordoni C, Chiarelli N, Ritelli M. Differential diagnosis and diagnostic flow chart of joint hypermobility syndrome/ehlers-danlos syndrome hypermobility type compared to other heritable connective tissue disorders. *Am J Med Genet C Semin Med Genet*. 2015; 169C (1): 6-22.
- [14] Beighton P, Solomon L, Soskolne CL. Articular mobility in an African population. *Ann Rheum Dis*. 1973; 32: 413-8.

- [15] Malek S, Reinhold EJ, Pearce GS. The Beighton Score as a measure of generalised joint hypermobility. *Rheumatol Int*. 2021; 41 (10): 1707-1716.
- [16] Grahame R, Bird HA, Child A. The revised (Brighton 1998) criteria for the diagnosis of benign joint hypermobility syndrome (BJHS). *J Rheumatol*. 2000; 27 (7): 1777-1779.
- [17] Hakim AJ, Grahame R. A simple questionnaire to detect hypermobility: an adjunct to the assessment of patients with diffuse musculoskeletal pain. *Int J Clin Pract*. 2003; 57 (3): 163-166.
- [18] De Wandele I, Rombaut L, Leybaert L, et al. Dysautonomia and its underlying mechanisms in the hypermobility type of Ehlers-Danlos syndrome. *Semin Arthritis Rheum*. 2014; 44 (1): 93-100.
- [19] Gazit Y, Nahir AM, Grahame R, Jacob G. Dysautonomia in the joint hypermobility syndrome. *Am J Med*. 2003; 115 (1): 33-40.
- [20] Botrus G, Baker O, Borrego E, et al. Spectrum of Gastrointestinal Manifestations in Joint Hypermobility Syndromes. *Am J Med Sci*. 2018; 355 (6): 573-580.
- [21] Castori M, Morlino S, Pascolini G, Blundo C, Grammatico P. Gastrointestinal and nutritional issues in joint hypermobility syndrome/Ehlers-Danlos syndrome, hypermobility type. *Am J Med Genet C Semin Med Genet*. 2015; 169C (1): 54-75.
- [22] Baeza-Velasco C, Bourdon C, Montalescot L, et al. Low- and high-anxious hypermobile Ehlers-Danlos syndrome patients: comparison of psychosocial and health variables. *Rheumatol Int*. 2018; 38 (5): 871-878.
- [23] Sanches SH, Osório Fde L, Uchina M, Martín-Santos R, Crippa JA. Anxiety and joint hypermobility association: a systematic review. *Braz J Psychiatry*. 2012; 34 Suppl 1: S53-S60.
- [24] Cederlöf M, Larsson H, Lichtenstein P, Almqvist C, Serlachius E, Ludvigsson JF. Nationwide population-based cohort study of psychiatric disorders in individuals with Ehlers-Danlos syndrome or hypermobility syndrome and their siblings. *BMC Psychiatry*. 2016; 16: 207.
- [25] Bulbena A, Baeza-Velasco C, Bulbena-Cabré A, et al. Psychiatric and psychological aspects in the Ehlers-Danlos syndromes [published correction appears in Am J Med Genet A. 2017 Dec; 173 (12): 3241]. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 237-245.
- [26] Hakim A, De Wandele I, O'Callaghan C, Pocinki A, Rowe P. Chronic fatigue in Ehlers-Danlos syndrome-Hypermobility type. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 175-180.
- [27] Krahe AM, Adams RD, Nicholson LL. Features that exacerbate fatigue severity in joint hypermobility syndrome/Ehlers-Danlos syndrome - hypermobility type. *Disabil Rehabil*. 2018; 40 (17): 1989-1996.
- [28] Henderson FC Sr, Austin C, Benzel E, et al. Neurological and spinal manifestations of the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 195-211.
- [29] Martin VT, Neilson D. Joint hypermobility and headache: the glue that binds the two together--part 2. *Headache*. 2014; 54 (8): 1403-1411.
- [30] Bendik EM, Tinkle BT, Al-shuik E, et al. Joint hypermobility syndrome: A common clinical disorder associated with migraine in women. *Cephalalgia*. 2011; 31 (5): 603-613.
- [31] Russek LN, Stott P, Simmonds J. Recognizing and Effectively Managing Hypermobility-Related Conditions. *Phys Ther*. 2019; 99 (9): 1189-1200.
- [32] Palmer S, Terry R, Rimes KA, Clark C, Simmonds J, Horwood J. Physiotherapy management of joint hypermobility syndrome--a focus group study of patient and health professional perspectives. *Physiotherapy*. 2016; 102 (1): 93-102.
- [33] Engelbert RH, Juul-Kristensen B, Pacey V, et al. The evidence-based rationale for physical therapy treatment of children, adolescents, and adults diagnosed with joint hypermobility syndrome/hypermobility Ehlers Danlos syndrome. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 158-167.
- [34] Kemp S, Roberts I, Gamble C, et al. A randomized comparative trial of generalized vs targeted physiotherapy in the management of childhood hypermobility. *Rheumatology (Oxford)*. 2010; 49 (2): 315-325.
- [35] Pacey V, Tofts L, Adams RD, Munns CF, Nicholson LL. Exercise in children with joint hypermobility syndrome and knee pain: a randomised controlled trial comparing exercise into hypermobile versus neutral knee extension. *Pediatr Rheumatol Online J*. 2013; 11 (1): 30.
- [36] Chopra P, Tinkle B, Hamonet C, et al. Pain management in the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet*. 2017; 175 (1): 212-219.