

Effect of Eucalyptus Globulus Woodlot Plantation on Selected Soil Physico-Chemical Properties, Wheat Yield in Wogera District, Amhara Region, Ethiopia

Kassegn Asnakew Seyoum¹, Bekalu Melis Alehegn¹, Asmamaw Alemu Abtew²

¹College of Agriculture and Natural Resource, Assosa University, Assosa, Ethiopia

²Collage of Agriculture and Environment Science, Gondar University, Gondar, Ethiopia

Email address:

Kassu367@gmail.com (K. A. Seyoum), bekalumelis6@gmail.com (B. M. Alehegn)

To cite this article:

Kassegn Asnakew Seyoum, Bekalu Melis Alehegn, Asmamaw Alemu Abtew. Effect of Eucalyptus Globulus Woodlot Plantation on Selected Soil Physico-Chemical Properties, Wheat Yield in Wogera District, Amhara Region, Ethiopia. *American Journal of Science, Engineering and Technology*. Vol. 6, No. 2, 2021, pp. 34-44. doi: 10.11648/j.ajset.20210602.14

Received: December 14, 2020; **Accepted:** December 25, 2020; **Published:** June 9, 2021

Abstract: Currently, a Eucalyptus globule is a common smallholder plantation species in the Wogera district. The species is significantly expanding on marginal and fertile farmlands. The objective is to analyze the effects of Eucalyptus globules woodlots plantation and their directions on selected soil physicochemical properties and yield of wheat crop. Three woodlots with similar site conditions were selected for this particular study. Soil samples were taken at a soil depth of 20 cm in RCBD design with three replications. Soil samples were taken at four directions (East, West, South, and North) of woodlots plantation at five different horizontal distances from the woodlots; the center of the woodlot, 5 m, 10 m, 15 m, and 40 m. For the wheat yield study: quadrates of 1m² area at each distance within directions were laid for wheat grain yield measurement. The finding indicated that electric conductivity, available phosphorus, cation exchange capacity, soil texture, and exchangeable cations were not significantly different across the different directions and distances, whereas soil bulk density, pH, total nitrogen, organic matter were significantly different. This could be attributed to the high organic matter addition via litter-fall, root biomass, uptake, and return of nutrients from deeper soil profiles under the tree canopies. The biomass and grain yield of wheat were significantly different across distance and direction from the sampled woodlot in increasing trend in all directions, this study recommends Eucalyptus should planting on marginal lands with appropriate distance from croplands about 20m to minimize its effect on crop yields.

Keywords: Eucalyptus, Soil Properties, Wheat, Distance, Direction

1. Introduction

The Eucalyptus is the diverse genus of flowering plants in the globe. It belongs to the family Myrtaceae and comprises about 800 species [1]. Worldwide area coverage of Eucalyptus plantation was estimated at about 17.9 million hectares [2]. The species was introduced to East Africa in the late 19th century and by the early 1970s. Currently, Eucalyptus plantation covers about 12 million hectares overall in the tropical zone. In Ethiopia, Eucalyptus was introduced during the regime of Emperor Menilek II (1868-1907) in 1894/95 [3]. The purpose was to supply fuelwood and construction timber to the new and growing capital city, Addis Ababa. Currently, about 5 to 10 Eucalyptus species are

commonly established for multipurpose uses [4]. Eucalyptus globules species are commonly planted on wide areas of land previously cultivated to crop production. Currently, an estimated area of 500,000ha of the land is covered by Eucalyptus plantation in Ethiopia [5]. Among these, Eucalyptus globules plantations cover more than 100,000 ha that growing well at elevations ranging from 1400-3200 m.a.s.l. The major energetic factors that farmers to grown Eucalyptus are: increasing demand for wood products in the market, high-income generation, high rate of biomass production, easy to establish, adaptability on infertile soils. [6] Farmers' increased their concentration in Eucalyptus plantation which caused the conversion of croplands into Eucalyptus plantation [7] However, the unrestrained

expansion of Eucalyptus on productive croplands has a great alarm, due to the detrimental effects on soil productivity. Eucalyptus tree might cause crop yield loss through extract essential nutrients and compete with crops and impoverish the soil [8].

In recent years, the expansion of Eucalyptus globules plantation on and around the cropland is expanding from year to year in an alarming rate especially in northern Ethiopia highlands including the present study area. Increasing plantations would create competition among land area of crops and Eucalyptus globules woodlot plantations. Similar trends of replacement of cropland with Eucalyptus woodlots have been observed across Ethiopian highlands [4]. Reports showed that Eucalyptus trees change watercourses, soil physical and chemical properties, propensity to exhaust soil nutrients, and fertility eventually affects crop production and productivity [9]. However, the effects of Eucalyptus species on soil physical and chemical properties and exhaustion of the soil essential nutrients still remain unclear [10] and scientific reports are insufficient [6]. Therefore, additional documentation on the specific effects of Eucalyptus globules plantations in relation to crop yield and soil properties in the district is required. Thus, this study was undertaken with the

aim to analyze the effects of Eucalyptus globules plantation on soil selected properties and wheat crop yield performances that grown neighboring to plantations. The results of the study can effectively create awareness for the community concerning specific effects of Eucalyptus globules tree species on crop performance and soil properties.

2. Materials and Methods

2.1. Description of the Study Area

The study was conducted in Wogera district, geographically situated at 12°46'06.5"N and 37°37'26.5"E in North Gondar Zone of Amhara region (figure 1). It is situated about 32 km away from Gondar city and 213 km Northwest of Bahir Dar, the capital city of Amhara region. The District covers an estimated area of 1821km². According to the Amhara Bureau of Finance and Economic Development (BoFED) 2012) the District has a total population of 250,493 out of this 127,858 are males and 122,635 are females, from the total population 94% live in rural areas, the population density is 137 person/km².




Figure 1. Map of the study area in the northwest Amhara region, Ethiopia.

2.2. Climate and Soil Type

According to the agro-ecological classification of Ethiopia, the study area is characterized by three agro-climatic zones highland (56%), Semi-lowland (26%), lowland (14%), and 4% frost (Wogera District agricultural office, 2013). The mean annual average temperature ranges between 140C to 330C. The rainfall pattern is orographic type, stretching from May to September. Annual average rainfall ranges between 1000 and 1600 mm which varies along the geographical zone. The District has an average altitude of 2812 meters above sea level (Wogera District agricultural office, 2013). Cambisol is the dominant soil type in the area and actually considered fertile

soils [11]. They are deep, well-drained, red, tropical soils. However, the area is highly populated there is severe erosion in the area. Since the landscape has lost its natural vegetation, the soil of the surrounding hills is degraded with frequent rock outcrops.

2.3. Land Use and Land Cover

The major land uses of the study area include cultivated land, grazing land and plantation of exotic species mainly *Eucalyptus globulus*. The large portion of the district is intensively cultivated land (Almost 68% of the land is under crop cultivation during cropping season). After crop

harvesting, the land is opened up for cropping and grazing livestock. Apart from few trees around homesteads and farm borders, native trees are rarely seen in the landscape of the study area... The community faced a shortage of wood for construction and firewood, therefore most farmers adopt and expand *Eucalyptus* plantation on their land to alleviate the problems (Wogera District agricultural office, 2016).

2.4. Livelihood

The study area community mainly depends on mixed agriculture. Agriculture is characterized by crop oriented small-scale subsistence farming-system with a wide range of cereals, pulses crops grown for household consumption. Commercial farming mainly involves the growing of *Eucalyptus globulus* and some food crops such as chickpea. In spite of the fact that the diversified potential of crop productivity farmers has widely planted *Eucalyptus globulus* on croplands, due to its fast growth and requires less care. The main plantation practices are woodlot, cropland boundary plantation, and road borders plantations to fulfill the needs for fuel wood, construction and income generation.

2.5. Sampling and Data Collection Methods

A combination of method was applied for the collection

and analysis of data to meet the objectives of the present study. Thus the selected and applied methods discussed in three categories below: i) soil sampling and analysis techniques to evaluate the effect of selected physical and chemical properties and ii) the method used to evaluate the effect of the woodlots on crop yield along different directions and distances.

2.5.1. Sampling of *Eucalyptus Globulus* Woodlot Stands

The experiment was set on existing *Eucalyptus* woodlots plantation and their surrounding wheat crop... A reconnaissance survey was carried out through transect walk and then three *Eucalyptus globulus* woodlot stands that bordered by wheat crop were selected. The woodlots stand were selected purposely having nearly similar soil management practices, altitude, slope gradient and *Eucalyptus* stand characteristics such as height, and diameter at breast height, age (all were \approx 13-14 years age). Measurements of tree dimensions taken such as: diameter at breast height using diameter tape, tree height using clinometer and crown diameter at crown base using measuring tape. Slope gradient and elevation of the study area were measured by using clinometer and global positioning system respectively.

Table 1. Characteristic features of the sampled *Eucalyptus* woodlot stands.

Attributes	Sampled <i>Eucalyptus globulus</i> stands		
	Stand 1	Stand 2	Stand 3
Stand type (Coppice, standard crop)	2 times coppice	2 times coppice	2 times coppice
Age (years)	14	13	13
Average height (m)	12.5	9.5	11
Average DBH (cm)	20	16	18
Slope (%)	5.5	6.25	7
Location (m.a.s.l)	2955	2959	2961
Previous land use	Cropland	cropland	Cropland
Surrounding land uses	Wheat crop land	Wheat crop land	Wheat crop land
Tillage frequencies	3 times	3 times	3 times
Rate of fertilizers (kg/ha)	100 kg DAP 100 kg Urea	100 kg DAP 50 kg Urea	100 kg DAP 100 kg Urea
Soil texture	Loam	Clay loam	Clay loam
Soil color	Blue	Blue	blue
Distance from homestead	300 m	265 m	290 m

The plantations were established in 2005 and were harvested three times. The average area of the woodlots was 0.375 ha, ranging from 0.25 ha to 0.5 ha. All woodlots were owned by farmers and planted in irregular pattern. The stand rotation age ranged from 4 to 5 years. Stand ages were 13-14 years and stand densities ranged from 9,500 to 12,000 individuals tree per ha.

2.5.2. Experimental Design and Data Collection of Crop Yield

The test crop was wheat (*Triticum aestivum*), the most commonly grown crop variety within the study area. According to farmer's crop calendar, wheat crop was sown during 10 June to 25 June, 2016 and harvested at the end of December, 2016. Wheat crop was locally seeded 120 kg and fertilized with 100 kg ha⁻¹ DAP (Di-Ammonium phosphate) and 100 kg ha⁻¹ Urea (nitrogen fertilizer at planting time).

Most Farmers has practiced weeding three times. As described by development agents and local farmers, growers could harvest greater than 50 quintals (5 tons) per hectare.

Under farmers' crop production circumstances, the effects of *Eucalyptus globulus* woodlot plantation on the neighboring wheat crop performances was assessed in the 2015/2016 cropping season. The field experimental independent variable considered were the distances (at 5m, 10m, 15m and 40m) and direction (oriented north, south, east and west) of the *Eucalyptus globules* woodlot stand (figure 2). At harvesting time, a plot areas of 1m*1m (1m²) were marked at 5m, 10m, 15m and 40 m [12] far from the woodlot stand along the four directions (East, West, North and South) of the woodlot base. The 40m distance represented sampling point, which was assumed to be free from tree influence, and thus used as control. The experimental factors and levels

were four directions and four distances, were arranged in RCBD (Random Complete Block Design) with three

replications. Thus, a total of 48 (4 distance x 3 woodlot x 4 direction x 1 depth) samples were collected.




Figure 2. Schematic presentation of the field experiment design.

Samples of wheat yield that grown at each sampled plot at harvesting time were collected at each distance in the four directions from the stand. The sample were air-dried, threshed manually and then weighted the grain yield in grams in December, 2016. After the yield was determined, the total dry biomass yield was also measured in a similar way.

2.6. Soil Sampling

Soil samples were collected in 2016 cropping season after wheat crops have been harvested. Similar design was followed for wheat yield sampling illustrated in Figure 2. Moreover, soil samples were taken at the center of the woodlot plantation. Soil samples were collected from four corners of each plot (1m²), with one from the center by auger from each laid distance and direction at the depth of 20cm. A total of 255 samples were collected from the three woodlot stands. Then the collected soil samples of sub-plot (at the four corners and center) of the plot were thoroughly mixed to form a composite sample. In this way, a total of 255 samples were converted into 51 composite soil samples for convenient laboratory analysis.

2.7. Soil Sample Preparation and Analytical Procedures

Before conducting the analysis, soil samples were air-dried, then grinded and sieved through the 0.5 and 2mm sieve to remove unwanted materials (crop/plant residue, gravels) from the sample except that of the bulk density sample. The soil samples were analyzed for chemical and textural properties at Gondar Soil Testing Laboratory using the following standard laboratory procedures [13]. Organic matter and total nitrogen contents of the soil were determined following the wet combustion method of the Walkley and Black method, and the wet digestion procedure of Kjeldhal

method, respectively [14]. Available P was extracted by the Olsen method [15]. PH and Electric conductivity (EC) (1:2.5, soil: water) of the soil were measured in water using a pH meter with a glass-calomel combination electrode [16]. Cation exchange capacity was determined following the 1nitrogen ammonium acetate method at pH 7 [17], and exchangeable calcium and magnesium were determined using EDTA (Ethylene diamine-tetra acetic acid) titration, whereas exchangeable potassium and sodium were determined from the same extraction with flame photometry. Soil texture was determined by using bouyoucos hydrometer method. An undisturbed core sample from the topsoil was taken to determine the bulk density [18]. Soil samples were oven-dried at 105oC for bulk density determination.

2.8. Data Analysis

Different analytical procedures were followed for the analysis of the two sets of crop yield and soil properties data. Analysis of variance using SAS (version 9) was performed to determine effects of *Eucalyptus globules* woodlot plantation on parameters of both crops and soil properties to the entire woodlot direction and distance. All analyses were performed at a probability level of 0.05, *i.e.*, statistical differences were determined by two-way ANOVA employing a 95% level of confidence. Descriptive statistical procedures were also applied.

3. Results and Discussion

3.1. Effect of *Eucalyptus Globulus* Plantation on Grain and Biomass Yield of Wheat

Eucalyptus globules plantations were assessed for their effect on the above ground biomass and grain yield of wheat

crop. The main effects (direction and distance) had highly significant effect on the grain and biomass yield of wheat ($P < 0.001$) (Table 2). The grain and biomass yield of wheat increased significantly as the distance (10 m) from the *Eucalyptus* stands increased in all directions (Figure 1).



Figure 3. Morning shading effect of *Eucalyptus globulus* plantation on crop lands.

The present findings concurred with that of [19] who concluded that the yield of wheat grain and straw increased significantly with an increase in distance from the stand. The increase in grain yield away from the woodlot could be due to less competition of nutrients, moisture, and shading effect under the woodlot canopies than the open fields. The maximum yield of wheat grain and biomass were 4893 and 5831 kg/ha respectively, recorded at the distance of 40m away from the woodlot at the Eastern direction, while the minimum yield of grain and above ground-biomass were 2309 and 3772 kg/ha respectively recorded at the distance of 5 m away from the woodlot at the western direction (Table 2). Consistent with the present findings, [20] also reported that the adjacent tree decreased maize height and biomass by 42.9% and 74.6% respectively in Tanzania. Similarly [21] also reported that 40% of crop yield reductions when crops close to trees in Kenya.

Table 2. Main effect of direction and distance on wheat crop yield.

Components	Wheat yield		
	Grain yield	Biomass/straw/ yield	
Directions	East	438.58a	528.19a
	West	335.27b	447.40b
	South	412.55c	500.36c
	North	409.73c	502.40c
	LSD	51.22	43.60
	Significant		
Distances (m)	5	335.71a	437.57a
	10	370.24a	472.50b
	15	411.27b	503.53c
	40	478.91c	564.75d
	LSD	40.24	30.40
	Significant		

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$.

The variation in grain and biomass yield between northern and southern directions was non-significant at respective woodlot plantation distance and direction. However, the grain and biomass yield between Eastern and Western directions varied significantly at all distances except 40m (Table 3). The maximum crop and biomass yield was observed on the Eastern side of woodlot stand base.

Table 3. Effect of *Eucalyptus* plantation on wheat grain and biomass yield (kg/ha).

Parameters	Directions	Distance (m) from the woodlot			
		5	10	15	40
Grain yield	East	3927a	4200b	4521c	4893d
	West	2309b	2644c	3813d	4643e
	North	3600a	3873b	4109c	4805e
	South	3591a	4091c	4006c	4813e
	CV (%)		2.69		
	LSD 0.05		17.85		
Biomass/straw/Yield	East	4730a	5086b	5480c	5831d
	West	3772b	4032bc	4502d	5588ed
	North	4443a	4931b	5164b	5556d
	South	4556a	4850b	4327c	5613d
	CV (%)		3.16		
	LSD 0.05		26.0		

Note: Column refers along direction and a row refers distance with the same letters are not significantly different at $p < 0.05$. The first letter represents significance comparison between distances and b represent between directions.

The interaction effect between distance and directions was significant (Table 3). There was a decline of nearly 35.3 and 52.8 percent of wheat grain and biomass yield respectively while comparing 5m and 40m intervals from the woodlot (Table 3). The effects of all directions at a 40 m distance were non-significant. These results are supported by other works, such as [22] that the high yield of wheat grain in areas without trees was presumably because of less competition of soil moisture, nutrients, and sunlight between agricultural crops and trees. They [23] studied the yield potential of inter-crop with *Eucalyptus* and its shading effect on adjoining wheat crops. The orientation of the tree line and the distance on which the associated crop they have grown contributed significantly towards crop growth and yield.

3.2. Effect of *Eucalyptus Globulus* Plantation on the Physical Properties

3.2.1. Soil Texture

According to the USDA textural classification, the texture of the area was clay loam soil. The sand proportions were (36.29%), silt (34.67%) and clay (29.04%) (Table 4). the soil of the areas was predominantly sand ($> 36.29\%$). The clay content in the soil was small (29.04%). The mean silt proportion of the soil under the woodlot center was lower than the open farmland (Table 4). Soil texture might affect soil physical and chemical and to some extent soil biological properties, was not significantly affected (Table 4) across distance and direction from woodlot the texture is more related to parent material than tree influence. According to Landon [24], clay, clay-loam, and silt-loam soils show low bulk density values (1.0 to 1.6 g cm⁻³) as compared to sands and sandy loam soils which show high bulk density values (1.2 to 1.8 g cm⁻³). The non-significant differences in the mean proportions of sand, silt, and clay fractions between the soils under the *Eucalyptus* woodlot and in the farmland suggests that the soils, are texturally similar, being clay loam and have derived from the same parent material, under the same climate, and similar topography.

Table 4. The main effect of Eucalyptus plantation direction and distance on soil texture and bulk density.

		Soil texture			Bulk density
		Sand	Clay	Silt	
Directions	East	37.94a	27.52a	34.53a	1.30a
	West	34.61a	30.58a	34.80a	1.21a
	South	36.44a	28.58a	34.97a	1.25a
	North	35.78a	28.58a	35.64a	1.27a
	LSD	5.67	6.36	5.74	0.11
	Significant	Ns	Ns	Ns	Ns
Distances (m)	Center (0)	37.94a	32.40a	29.65a	1.13a
	5	34.78a	30.80a	34.41a	1.19a
	10	35.44a	28.74a	35.80a	1.25a
	15	37.11a	26.52a	36.36a	1.26a
	40	37.44a	29.18a	33.36a	1.34a
	LSD	8.61	9.11	6.02	0.12
	Significant	Ns	Ns	Ns	Ns

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$.

3.2.2. Bulk Density

The bulk density of the soils ranged from 1.13 (center of woodlot) to 1.34 g cm⁻³ (Table 4), which categorized as low [25]. Similar findings, [26] reported lower bulk density under Eucalyptus plantation which ranges 1.07 for topsoil for clay soil in the Amhara Regional State of Ethiopia. [27] Reported a lower bulk density under Eucalyptus plantation and Croton macrostachyus ranged 1.0 to 1.1 g/cm⁻³ for topsoil (0-20 cm) in loamy clay soil in Lake Tana plain of the same region.

The soil bulk density increased from 1.13 to 1.34 g/cm³ from the woodlot center to wards adjacent farmland at all directions (Table 4). However, there was no significant difference of bulk density among the distance from the woodlot of the *Eucalyptus* plantation. Meanwhile, there was not significantly differing of bulk density along the directions and interaction effect. However, the highest bulk density of (1.34 g cm⁻³) was observed at 40m distance in the north direction. The lower bulk density was recorded in east direction. On contrary, [28] reported that lower bulk density in crop lands as compared to *Eucalyptus* plantation. The lower bulk density might be due to organic matter accumulation and microbial activity.

Table 5. The effect of Eucalyptus plantation on soil pH at different distance and direction.

Parameter	Directions	Distance (m) from the woodlot				
		Center	5	10	15	40
pH	East	5.28a	5.68b	5.84b	5.67b	5.59b
	West	5.28a	5.69b	5.58b	5.63b	5.68b
	North	5.28a	5.50b	5.54b	5.39b	5.37b
	South	5.28a	5.79b	5.67b	5.45b	5.51b
	CV (%)	5.01				
	LSD 0.05	0.38				

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$; the first letter represents significance comparison between distances and b represent between directions.

3.3. Effects of Eucalyptus Globulus Plantation on Soil Chemical Properties

3.3.1. Soil pH

The pH values of the samples taken from surface layer were ranged from 5.28 to 5.84 (Table 5). Soil pH was not significantly different on the main and interaction effects of the woodlot stand. Generally, lower mean soil pH values were recorded at the center of woodlot as compared to soils away from the woodlot which might be due to several mechanisms that release H⁺ ions, such as soil base cation uptake by the tree, decomposition of organic matter to organic acids and CO₂, root respiration and nitrification.

According to Tekalign *et al.* rating of soil pH, soils with pH (H₂O) > 8.0 are characterized as strongly alkaline [29]; 7.4-8.0 as moderately alkaline; 6.7-7.3 as neutral, while soils with pH of 6.0-6.6, 5.3-5.9, 4.5-5.2 and < 4.5 are rated as slightly acid, moderately acid, strongly acid and very strongly acid, respectively. According to the study result the soil was moderately acidic, that ranges between 5.7 and 6.8, which is good for crop production as most nutrients for field crops are available at pH values of greater than 5.5 [24]. These findings supported by others; lower soil pH recorded under *Eucalyptus* woodlot plantation as compared to farmlands and grazing lands in Ethiopian central highlands [24]. Zewdie also found that lower pH under *Eucalyptus globulus* plantations than agricultural lands [30].

3.3.2. Soil Organic Matter

Soil organic matter was ranged from 2.32 to 6.42 (Table 7) classified as high Soil organic matter level [31] who classified organic carbon as low (0.6-1.16%), moderate (1.16-1.74%) and high (>1.74%). There was significantly different in soil organic matter between the distance the woodlot and directions ($P < 0.05$). The interaction effect between distance from the woodlot and directions was not significant (Table 7). The soil organic matter were higher at the center of the woodlot than away the woodlot of the nearby cropland that showed a reducing trend with increasing distance from the center of woodlot towards the crop field with in all directions.

Table 6. The main effect of *Eucalyptus* plantation direction and distance on soil chemical properties.

		Soil chemical properties								
		pH	EC	OM	TN	CEC	Av. P	Ex. Ca	Ex. Mg	Ex. K
Directions	East	5.69a	0.027a	3.16a	0.16a	32.81a	25.61a	17.43a	13.63a	1.52a
	West	5.64a	0.024a	2.45a	0.12a	33.15a	15.17a	16.96a	13.97a	1.54a
	South	5.61a	0.025a	3.03a	0.14a	33.73a	24.41a	17.24a	14.63a	1.50a
	North	5.45a	0.028a	3.04a	0.15a	31.77a	17.39a	14.62b	14.88a	1.42a
	LSD	0.25	0.010	1.22	0.06	5.53	15.29	2.12	5.26	0.27
Sig.										
Distances (m)	Center	5.28a	0.030a	6.41a	0.32a	35.73a	8.41a	14.90a	18.07a	1.44a
	5	5.66b	0.025a	3.26b	0.16b	34.76a	21.10b	17.71a	14.85a	1.58a
	10	5.65b	0.030a	2.97b	0.14b	34.57a	23.80b	16.87a	13.26a	1.59a
	15	5.54b	0.025a	2.64b	0.13b	32.03a	19.97b	15.39a	13.12a	1.39a
	40	5.40b	0.024a	2.82b	0.13b	30.10a	17.71b	16.28a	15.88a	1.43a
LSD		0.34	0.014	1.70	0.08	5.41	22.68	3.53	6.24	0.33
Sig.										

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$; 'a' letter represents significance comparison between distances and 'b' represent between directions.

Soil organic matter content at eastern direction was high whereas less content at western direction were recorded than other directions (Table 7), thus might be passive microbial activities because of late sunlight coming due to shading in the western. The soil organic matter of woodlot center was 2.76 times higher than the respective soil surface of control (40m away from the woodlot stand). This variation in soil

organic matter with distance away from the woodlot center could be due to accumulation of leaf litters, seeds fall and dead roots from the tree as compared to the adjacent crop lands while the source of soil organic matter outside the woodlot was mostly crop residues which might be removed as a fuel wood and also grazing.

Table 7. The effect of *Eucalyptus* plantation on soil organic matter and total nitrogen at different distance and direction.

Parameters	Directions	Distance (m) from the woodlot				
		Center	5m	10m	15m	40m
Soil organic matter (%)	East	6.42a	3.50b	3.45b	2.82b	2.89b
	West	6.42a	2.46b	2.47b	2.56b	2.33b
	North	6.42a	3.52b	3.01b	2.73b	3.27b
	South	6.42a	3.57b	2.96b	2.82b	2.79b
	CV (%)		38.6			
	LSD 0.05		2.9			
Soil total nitrogen (%)	East	0.32a	0.17b	0.17b	0.14b	0.14b
	West	0.32a	0.12b	0.12b	0.12b	0.11b
	North	0.32a	0.18b	0.15b	0.12b	0.16b
	South	0.32a	0.18b	0.15b	0.14b	0.14b
	CV (%)		40.3			
	LSD 0.05		0.14			

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$; 'a' letter represents significance comparison between distances and 'b' represent between directions

3.3.3. Soil Total Nitrogen

The total nitrogen contents of the soils were ranged from 0.11 to 0.32% (Table 7), which could be categorized as medium to high [32]. Who categorized soil total nitrogen rate as very low (<0.1), low (0.1-0.15) medium (0.15-0.25), and high (>0.25). Total nitrogen was significantly different among directions from the woodlot stand (Table 7). In the western direction lower total nitrogen was observed. However, soil nitrogen was not significantly affected by the distance and directions from woodlot. Total nitrogen was significantly higher at the center of the woodlot than the adjacent farmland and this might be attributed to more organic matter and nitrogen mineralization at the center of *Eucalyptus* plantation. The other reason for high soil nitrogen under *Eucalyptus* woodlot might be due to the low temperature and very limited radiation reached on the surface

of the soil that leads to low volatilization of $\text{NH}_3\text{-N}$ [33].

There was slightly decrease in total nitrogen with distances, higher at Eastern direction as compared to other directions. Soil nitrogen concentration showed decreasing trend with increasing distance from woodlot plantation within all directions (Table 7), this could be due to the effect of intensive and continuous cultivation forced oxidation of organic carbon of field that release nitrogen from organic matter decomposition [34].

The deep root structure of *Eucalyptus* can operate as a nutrient pump for leached nitrogen from deep soil or slow down leaching. Soils under *Eucalyptus* plantation reported as it has larger soil nitrogen due to the long time under tree cover and soil nitrogen mineralization could be increased under *Eucalyptus* plantation by 11-14 ppm per year [26]. He stated that total nitrogen increased with distance from the

Eucalyptus trees, their result showed that near the *Eucalyptus* stand the soil total nitrogen was significantly larger than the average soil total nitrogen of the field and the nitrogen content declines with increasing distance up to 5 m away from the woodlot plantation (Table 7). Generally, higher soil nitrogen near *Eucalyptus* trees might be due to the higher nitrogen in the foliage of *Eucalyptus tree* and low uptake by the tree from the top soil.

3.3.4. Available Soil Phosphorus

The available phosphorus contents of the soils were ranged from 8.42 to 30.31 ppm (Table 8) that could be medium to high phosphorus, according to Josan et al. [35] Who categorized as very low (<3), low (4-7), medium (8-11), and

high (>12). Soil phosphorus concentration was not affected by the main effects of distance from the woodlot and directions and their interaction effect (Table 8). Soil phosphorus concentration increased with increasing distance from the woodlot with in all directions. The mean soil phosphorus concentration at the center of woodlot was 8.42 ppm lower than the nearby cropland. The lower soil phosphorus accumulation was observed at the center of the woodlot zone than adjacent crop land (Table 8) this could be due to fertilizer application on the adjacent farmland annually. The phosphorus concentration at western direction was lower towards the cropland.

Table 8. The effect of *Eucalyptus* plantation on available phosphorus, cation exchange capacity and electric conductivity at different distance and direction.

Parameters	Directions	Distance (m) from the woodlot				
		Center	5m	10m	15	40m
Available phosphorus (ppm)	East	8.42a	21.50a	28.80a	30.31a	21.83a
	West	8.42a	18.70a	14.92a	12.62a	14.47a
	North	8.42a	18.16a	21.20a	17.02a	13.21a
	South	8.42a	26.04a	30.31a	19.93a	21.36a
	CV (%)	83.06				
	LSD 0.05	27.4				
Cation exchange capacity (Cmol/kg)	East	36.73a	33.96a	34.44a	29.19a	33.66a
	West	36.73a	35.31a	31.28a	30.64a	34.38a
	North	36.73a	32.13a	29.48a	28.11a	37.35a
	South	36.73a	35.67a	32.92a	32.45a	32.89a
	CV (%)	14.60				
	LSD 0.05	8.00				
Electric conductivity (ds/m)	East	0.03a	0.026a	0.03a	0.023a	0.023a
	West	0.03a	0.02ba	0.023ba	0.026ba	0.026ba
	North	0.03a	0.03ab	0.03a	0.026ab	0.023a
	South	0.03a	0.026ba	0.023a	0.023a	0.02a
	CV (%)	40.63				
	LSD 0.05	0.01				

Note: column refers along direction and rows refer distance with the same letters are not significantly different at $p < 0.05$; 'a' letter represents significance comparison between distances and 'b' represent between directions.

These findings supported by other works; the available phosphorus content of farmland is higher than forest lands, which suggests that the annual leftover of phosphorus from fertilizer application is more than obtained from leaf decomposition from trees [36, 26] also found that the amount of available phosphorus content in croplands is much larger than in forest and *Eucalyptus* woodlots. Bewket and Stroosnijder reported that lower available phosphorus content at forest lands is due to a high proportion of phosphorus which retained and immobilized by microbes [37]. Another possible reason for low available phosphorus under *Eucalyptus* woodlots plantation could be due to the lower pH conditions that can permanently fix phosphorus. Under acid conditions, phosphorus is precipitated as Fe or Al phosphates of low solubility [26], whereby maximum availability of phosphorus generally happens in range of 6.0 to 7.0 soil P^H .

3.3.5. Cation Exchange Capacity

The cation exchange capacity of the soils was not significant for their main and interaction effect between distance and direction from the woodlot, but the higher value

of cation exchange capacity was observed at the center of the *Eucalyptus* woodlot stand (Table 8), this could be depletion of exchangeable bases as a result of intensive cultivation and application of inorganic application fertilizers which reduced the cation exchange capacity. Generally cation exchange capacity decreased with increasing distance from the woodlot within all directions (Table 8). This could be mainly due to high of accumulation organic matter at the center of the woodlot than the nearby cropland. The higher amounts of soil organic matter under the tree *Eucalyptus* plantation may imply that more cations would be released to the soil through mineralization as a result; the amount of negative charges in the soil would be higher. Our result is confirmed that cation exchange capacity didn't show any significant difference among the land uses in the central highlands of Ethiopia [26].

3.3.6. Soil Electrical Conductivity

The analysis of variance for soil electrical conductivity revealed that they were not significantly affected by interaction effect and main effects of distance and directions (Table 8). This could probably be due to high rainfall amount that remove base forming cations from the surface soil. In

general, higher soil electrical conductivity at the center of the woodlot than the adjacent crop land, this might be due to the increased accumulation of above ground biomass and associated cation uptake by the tree. [38] Suggested that soils with soil electrical conductivity value of below 0.80 dS m^{-1} are considered normal and suitable for all crop types. The soil electrical conductivity soil at the center of woodlot was 0.03 dS m^{-1} while that of beyond woodlot ranged from 0.03 to 0.02 dS m^{-1} (Table 8). Hence, the soils of the adjacent to crop land are suitable for most crops. According to [39] in the soil, the determination of the electrical conductivity serves to give an idea of the total quantity of soluble salts and the degree of salinity. The critical level of electrical conductivity of saturated soil paste extracts (ECe) for most crops is 4 dS m^{-1} . Soils with ECe greater than 4 dS m^{-1} are saline soils. These are considered restrictive for most crops and values as low as 2 dS m^{-1} may affect the more sensitive crops.

3.3.7. Soil Exchangeable Cations

The values of three primary exchangeable cations (calcium, magnesium and potassium) were not significantly different within distance, direction and interaction effect between distances and direction from woodlot, but gradually level of exchangeable cations decreasing with increasing distance from the woodlot at all directions (Table 8). There was higher exchangeable magnesium at the center of the woodlot plantation (Table 8), which could be due to the high accumulation of litter under plantation and decrease their reduction from leaching or erosion. But lower level of exchangeable calcium and potassium were observed at the of the *Eucalyptus* woodlot stand (Table 8). Findings supported by [40] who found lower level of calcium, magnesium and potassium under *Eucalyptus* tree at 0-20 cm soil depth.

The result shows that lower exchangeable calcium and potassium cations amount under plantation; this indicated that the *Eucalyptus* trees absorb these nutrients in standing biomass faster than recycle i.e. nutrients back to the soil. The immobilization of exchangeable bases, especially calcium resulted in lower soil pH under *Eucalyptus* plantation at 0–20 cm soil depth [41]. According to [42], the prevalence of calcium followed by magnesium and potassium in the exchange site of soils is favorable for crop production.

4. Conclusion

The experimental result showed that *Eucalyptus* tree and its direction significantly affect crop yield. The crop yield shown that increasing progressively as the distance from *Eucalyptus* woodlot stand base increases at all directions, hence higher yields were recorded in eastern direction whereas lower yields were found at the western direction. This work revealed that there was around 14.2 fold wheat grain yield difference from the stand as compared with the control plot (40m) and so did for biomass yield. The *Eucalyptus* woodlot and its direction have a considerable effect, in that western direction is more pronounced. This study indicated that under *Eucalyptus* woodlot plantation,

soils did not vary significantly in texture cation exchange capacity, electric conductivity, potassium, available phosphors, exchangeable calcium, magnesium and potassium cations and electric conductivity, on the other hand soil properties such as organic matter, total nitrogen, soil pH, and bulk density were significantly higher at the center of *Eucalyptus* woodlot plantation than in the adjacent cropland with decreasing trend with distance from the woodlot stand in all directions. Soil organic matter status under *Eucalyptus globules* plantation stand is higher, this may be leaf and root litter addition. Higher values of organic matter, total nitrogen and available phosphors were found in eastern direction, but lower in western direction. Overall, the studied results lead that *Eucalyptus globules* plantation affects wheat crop yield and selected soil physical and chemical properties. To minimize the adverse effects and improve the environmental importance of *Eucalyptus* tree plantation, choice of suitable site and management methods is very important. Moreover, when grown on cropland, *Eucalyptus* plantation should be limited to appropriate sites with right management and utilization so that it will not unfavorably affect neighboring crop productivity.

5. Recommendation

For the sustainable crop production, *Eucalyptus* tree should be planted in the order of west, north, south and east direction of the cropland. The result of wheat yields reported in this study were under farmer's crop production and management practices which may not be applied the same management practice. So, further study is needed under controlled experiment in association with this *Eucalyptus* plantation. Hence, further research required on effect of its leave for soil physical and chemical properties. Moreover, soil microbial population and their activities with *Eucalyptus* plantation aspects should be studied in details.

References

- [1] Adhanom, Daniel, & Toshome, Tefera. (2016). Characterization and classification of soils of Aba-Midan sub watershed in Bambasi Wereda, West Ethiopia. *International Journal of Scientific and Research Publications*, 6 (6), 390-399.
- [2] Alebachew, Melkamu, Dar, Bahir, Dar, Bahir, & Dar, Bahir. (2015). Investigation of the Effects of *Eucalyptus camaldulensis* on Performance of Neighbouring Crop Productivity in Western Amhara, Ethiopia. *Open Access Library Journal*, 2 (03), 1.
- [3] Alemie, Tilashwork Chanie. (2009). *The effect of eucalyptus on crop productivity, and soil properties in the Koga Watershed, Western Amhara Region, Ethiopia*. Cornell University.
- [4] Atumo, Tessema Tesfaye. (2018). Evaluation of Forage Type Cowpea (*Vigna unguiculata* L. Walp.) Accessions for Dry Matter Yield in Lowlands of Southern Ethiopia. *Evaluation*, 66.

- [5] Berhe, Daniel Hagos, Anjulo, Agena, Abdelkadir, Abdu, & Edwards, Sue. (2013). Evaluation of the effect of *Ficus thonningii* (blume) on soil physicochemical properties in Ahferom district of Tigray, Ethiopia. *Journal of Soil Science and Environmental Management*, 4 (2), 35-45.
- [6] Bewket, Woldeamlak, & Stroosnijder, Leo. (2003). Effects of agroecological land use succession on soil properties in Chemoga watershed, Blue Nile basin, Ethiopia. *Geoderma*, 111 (1-2), 85-98.
- [7] Blake, G R, & Hartge, KH. (1986). Bulk density. *Methods of soil analysis: Part 1 Physical and mineralogical methods*, 5, 363-375.
- [8] Buraka, Tadele, Sorsa, Zemach, & Lelago, Alemu. (2016). Response of Faba Bean (*Vicia Faba L.*) To phosphorus fertilizer and farm yard manure on acidic soils in Boloso sore Woreda, Wolaita Zone, Southern Ethiopia. *Food Science and Quality Management*, 53, 2225-0557.
- [9] Chanie, Tilashwork, Collick, Amy S, Adgo, Enyew, Lehmann, C Johannes, & Steenhuis, Tammo S. (2013). Eco-hydrological impacts of Eucalyptus in the semi humid Ethiopian Highlands: the Lake Tana Plain. *Journal of Hydrology and Hydromechanics*, 61 (1), 21-29b.
- [10] Chapman, HD. (1965). Cation-exchange capacity. *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*, 9, 891-901.
- [11] Dessie, Abebe Birara, Abteu, Asmamaw Alemu, & Koye, Abebe Dagnew. (2019). Determinants of the production and commercial values of Eucalyptus woodlot products in Wogera District, Northern Ethiopia. *Environmental Systems Research*, 8 (1), 4.
- [12] Egata, Desta Fikadu, & Desta, Wondimkun Dikir. (2020). Determination of Optimum Spacing and Harvesting Age of Citron Scented Gum (*Eucalyptus citriodora H.*) For Maximum Biomass and Essential Oil Yield at Wondo Genet, South Ethiopia.
- [13] El-Amin, EA, Diab, IE, & Ibrahim, S. (2001). Influence of Eucalyptus cover on some physical and chemical properties of a soil in Sudan. *Communications in soil science and plant analysis*, 32 (13-14), 2267-2278.
- [14] Gindaba, Jiregna, Rozanov, Andrey, & Negash, Legesse. (2007). Depletion of nutrients in adjacent crop lands by *Eucalyptus camaldulensis*. *South African Journal of Plant and Soil*, 24 (1), 47-50.
- [15] Gizachew, Kebede. (2017). Expansion of eucalypt woodlot and its factors in Cheha District, Southern Ethiopia. *World Scientific News*, 66, 163-180.
- [16] Jaleta, Daniel, Mbilinyi, Boniface, Mahoo, Henry, & Lemenih, Mulugeta. (2016). Eucalyptus expansion as relieving and provocative tree in Ethiopia. *Journal of Agriculture and Ecology Research International*, 1-12.
- [17] Jenbere, Dereje, Lemenih, Mulugeta, & Kassa, Habtemariam. (2012). Expansion of eucalypt farm forestry and its determinants in Arsi Negelle District, South Central Ethiopia. *Small-Scale Forestry*, 11 (3), 389-405.
- [18] Josan, MS, Nair, VD, Harris, WG, & Herrera, D. (2005). Associated release of magnesium and phosphorus from active and abandoned dairy soils. *Journal of environmental quality*, 34 (1), 184-191.
- [19] Kebede, YECHALE, & Raju, SAJ. (2011). Effect of land use/land cover change on soil properties in the Hare River Watershed, Ethiopia. *The Ecoscan, An International Quarterly Journal of Environmental Sciences*, 5, 69-74.
- [20] Khaliq, Abdul, & Kharr, Raja Attaullah. effect of eucalyptus *camaldulensis* on the yield of wheat and maize crops after reducing tree density.
- [21] Landon, John Richard. (2014). *Booker tropical soil manual: a handbook for soil survey and agricultural land evaluation in the tropics and subtropics*: Routledge.
- [22] Liang, Janice, Reynolds, Travis, Wassie, Alemayehu, Collins, Cathy, & Wubalem, Atalel. (2016). Effects of exotic *Eucalyptus* spp. Plantations on soil properties in and around sacred natural sites in the northern Ethiopian Highlands. *AIMS Agriculture and Food*, 1 (2), 175-193.
- [23] Mengist, Muluneh, Georg, G, & Sieghardt, M. (2011). Eucalyptus plantations in the highlands of Ethiopia revisited: A comparison of soil nutrient status after the first coppicing. *Mountain Forestry Master Programme*.
- [24] Mensah, ALBERT KOBINA. (2016). *Effects of eucalyptus plantation on soil physico-chemical properties in Thiririka sub-catchment, Kiambu County, Kenya*. Kenyatta University.
- [25] Mouazen, Abdul Mounem, Karoui, R, De Baerdemaeker, Josse, & Ramon, Herman. (2005). Classification of soil texture classes by using soil visual near infrared spectroscopy and factorial discriminant analysis techniques. *Journal of near infrared spectroscopy*, 13 (4), 231-240.
- [26] Mugayi, Emmanuel. (2019). *Impacts of Eucalyptus plantation on soil physico-chemical properties in Bamunanika sub-county, Luwero district, Uganda*. Makerere University.
- [27] Mulumba, Lukman Nagaya, & Lal, Rattan. (2008). Mulching effects on selected soil physical properties. *Soil and Tillage Research*, 98 (1), 106-111.
- [28] Nelson, Darrell W, & Sommers, Lee E. (1996). Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 3 Chemical methods*, 5, 961-1010.
- [29] Oballa, PO, Konuche, PKA, Michiri, MN, & Kigomo, BN. (2010). Facts on growing and use of Eucalyptus in Kenya.
- [30] Odhiambo, HO, Ong, CK, Deans, JD, Wilson, J, Khan, AAH, & Sprent, JI. (2001). Roots, soil water and crop yield: tree crop interactions in a semi-arid agroforestry system in Kenya. *Plant and soil*, 235 (2), 221-233.
- [31] Oehl, Fritz, Laczko, Endre, Bogenrieder, Arno, Stahr, Karl, Bösch, Robert, van der Heijden, Marcel, & Sieverding, Ewald. (2010). Soil type and land use intensity determine the composition of arbuscular mycorrhizal fungal communities. *Soil Biology and Biochemistry*, 42 (5), 724-738.
- [32] Olsen, Sterling Robertson. (1954). *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*: US Department of Agriculture.
- [33] Ralhan, PK, Singh, A, & Dhanda, RS. (1992). Performance of wheat as intercrop under poplar (*Populus deltoides* Bartr.) Plantations in Punjab (India). *Agroforestry systems*, 19 (3), 217-222.
- [34] Rayment, George E, & Lyons, David J. (2011). *Soil chemical methods: Australasia* (Vol. 3): CSIRO publishing.

- [35] Selassie, Yihenew G, & Ayanna, Getachew. (2013). Effects of different land use systems on selected physico-chemical properties of soils in Northwestern Ethiopia. *Journal of agricultural science*, 5 (4), 112.
- [36] Sharma, Asha, & Sah, vk. comparative growth, yield and yield attributes of wheat under poplar and eucalyptus based agroforestry system.
- [37] Sharu, MB, Yakubu, M, Noma, SS, & Tsafe, AI. (2013). Characterization and classification of soils on an agricultural landscape in Dingyadi District, Sokoto State, Nigeria. *Nigerian Journal of Basic and Applied Sciences*, 21 (2), 137-147.
- [38] Shirokova, Y, Forkutsa, I, & Sharafutdinova, N. (2000). Use of electrical conductivity instead of soluble salts for soil salinity monitoring in Central Asia. *Irrigation and Drainage Systems*, 14 (3), 199-206.
- [39] Tekalign, Tadesse, Haque, I, & Aduayi, EA. (1991). Soil, plant, water, fertilizer, animal manure and compost analysis. *Working document* (13).
- [40] Wedajo Abdi, Mulisa. (2019). *Localizing Ethiosis Fertility Map Based Fertilizer Type Recommendation for Maize (Zea mays L.) In Coffee and Spice Production Belt in Yeki District, Southwest of Ethiopia*. Jimma University.
- [41] Yitaferu, Birru, Abewa, Anteneh, & Amare, Tadele. (2013). Expansion of eucalyptus woodlots in the fertile soils of the highlands of Ethiopia: could it be a treat on future cropland use? *Journal of Agricultural Science*, 5 (8), 97.
- [42] Zewdie, Mulugeta. (2008). *Temporal changes of biomass production, soil properties and ground flora in Eucalyptus globulus plantations in the central highlands of Ethiopia* (Vol. 2008).
- [43] Zewdie, Mulugeta, Olsson, Mats, & Verwijst, Theo. (2009). Above-ground biomass production and allometric relations of Eucalyptus globulus Labill. Coppice plantations along a chronosequence in the central highlands of Ethiopia. *Biomass and Bioenergy*, 33 (3), 421-428.