

Antioxidant and Antiinflammatory Activities of *Imperata cylindrica* (L.) P. Beauv. (Poaceae) Extracts in Relation to Extraction Methods and Plant Organ

Anne Marie Leticia Konan^{1,*}, Esmel Essis Lohoues¹, Félix Adjé², Koffi Julien Golly³, Adou Koffi Matthieu Kra⁴, Kessé Phyllipe N'da², Amissa Augustin Adima²

¹Biochemical Laboratory, UFR Medical Sciences, Felix Houphouet-Boigny University, Abidjan, Côte d'Ivoire

²Research Department of Water Chemicals and Natural Substances, INP-HB, Yamoussoukro, Côte d'Ivoire

³Immunity Biology Center, Institut Pasteur, Abidjan, Côte d'Ivoire

⁴Pharmacodynamics Biochemical Laboratory, UFR Biosciences, Felix Houphouet-Boigny University, Abidjan, Côte d'Ivoire

Email address:

letitiakonan@hotmail.com (Anne Marie Leticia Konan), lohouese6@yahoo.fr (Esmel Essis Lohoues), felix_adje@yahoo.fr (Félix Adjé), gollykof@yahoo.fr (Koffi Julien Golly), mattykra@yahoo.fr (Adou Koffi Matthieu Kra), kphilipnda@gmail.com (Kessé Phyllipe N'da), amissaaug@gmail.com (Amissa Augustin Adima)

*Corresponding author

To cite this article:

Anne Marie Leticia Konan, Esmel Essis Lohoues, Félix Adjé, Koffi Julien Golly, Adou Koffi Matthieu Kra et al. (2023). Antioxidant and Antiinflammatory Activities of *Imperata cylindrica* (L.) P. Beauv. (Poaceae) Extracts in Relation to Extraction Methods and Plant Organ. *Biomedical Sciences*, 9(4), 97-103. <https://doi.org/10.11648/j.bs.20230904.13>

Received: December 6, 2023; **Accepted:** December 19, 2023; **Published:** December 28, 2023

Abstract: The involvement of oxidative stress/inflammation couple in the pathophysiology of many diseases increasingly emphasized. Treatments controlling the inflammatory process and the production or elimination of free radicals may be useful in limiting their harmful effects. *Imperata cylindrica* is a widely used medicinal plant in Côte d'Ivoire due to its numerous biological properties. This study investigated the antioxidant and anti-inflammatory activity of *Imperata cylindrica* organs in relation to different extraction methods. Leaves and roots of *I. cylindrica* collected from Côte d'Ivoire were washed, shade-dried and pulverized for storage. Phytochemical screening of water, alcohol and acetone raw extracts from roots and leaves was performed using conventional methods. Aqueous, alcoholic and acetonic decoctates and macerates from the leaves and roots of the plant prepared according to methodologies used in the literature were used to measure antioxidant and antiinflammatory activities by methods of inhibition of lipoperoxidation and denaturation respectively. The comparative study showed a difference in the extraction rate of bioactive compounds as well as the biological properties. The most suitable parameter for extracting antioxidant and anti-inflammatory compounds turned out to be the trio: roots / water / boiling. The richness of bioactive substances would justify the antioxidant and anti-inflammatory activities of the plant.

Keywords: Antioxidant, Antiinflammatory, Extraction, Leaves, Roots, *Imperata cylindrica*

1. Introduction

Inflammation results from the stimulation of the innate immune system by endogenous factors (IL33, IL- α , HMGB1, alarms) and exogenous factors (bacteria, viruses, allergens, toxic compounds, foreign bodies) in order to defend, restore and maintain cellular balance through a series of cellular and molecular reactions [1-3]. An inappropriate inflammatory response is the cause of loss of function, associated with

numerous pathologies (cardiovascular diseases, cancer, atherosclerosis, diabetes, hypertension and other degenerative diseases) [4-6]. Protein denaturation associated with inflammation is thought to occur when proteins have lost their structures and functions [8]. This denaturation causes conversion of phospholipids into eicosanoids and a mixture of oxidized phospholipids which can therefore be pro- and anti-inflammatory inducers causing an aggravation of the inflammatory process [8, 9, 11].

Free radicals are atoms or molecules resulting from metabolism and redox reactions that take place at multiple cellular levels [12]. Their excessive production, associated with the activity of proteolytic enzymes and hyperactivated leukocytes, is thought to be the cause of tissue and cellular damage [7]. Indeed, free radicals interact with proteins and lipids [14].

In order to avoid their harmful effects, the body has put in place defences (vitamins and enzymes) to neutralise them [13]. When these means are exceeded and an accumulation of free radicals is observed, a cascade of radical reactions is triggered leading to oxidative stress [14].

Lipoperoxidation is known to be the main consequence resulting of this state [15]. Membrane phospholipids and polyunsaturated fatty acids (mainly linoleic, linolenic, arachidonic and eicosapentaenoic acids) constituted the preferred targets of oxygenated free radicals and other reactive oxygen species [15-17]. Lipoperoxidation products are known to play a major role in the deregulation of key pathways in the pathophysiology of many diseases [18, 19].

Treatments able to controlling the inflammatory process and/or oxidative stress would be an alternative to limit the consequences of oxygenated free radicals [5, 13, 20].

Metabolites of medicinal plants are known to regulate and modulate various functions and processes, including inflammatory response and oxidative stress [21]. Their extractions constitute an important step in the discovery of bioactive components. Thus, biological activities of extracts have often presented significant differences depending on the extraction methods [22].

Imperata cylindrica is a *Poaceae* (*Monocotyledon, cyperales*) from tropical, subtropical or warm temperate regions [23]. In Côte d'Ivoire, it is found throughout the territory where it is used in the treatment of anemia and sinusitis. Furthermore, the plant is used alone or in combination with other plants, in decoction or maceration, for its antihypertensive, neuroprotective, antibacterial, antihelminthic, astringent, anti-inflammatory, digestive diuretic, emollient, febrifuge, hemostatic, antihyperglycemic, antidiarrheal and antioxidant properties. [25, 26, 28].

Phytochemical studies of the plant have highlighted presence of alkaloids, phenolic acids, flavonoids, tannins, glycosides, saponins, phenylpropanoids, sesquiterpenes [28-30].

This study investigated antioxidant and anti-inflammatory activity in relation to different methods of extraction of roots and leaves of *Imperata cylindrica*.

2. Materials and Methods

2.1. Material

Leaves and roots of *I. cylindrica* collected in central Côte d'Ivoire, were identified under number 051 of the Africa Rice herbarium (Rice Center for Africa, Bouaké). They were washed, then shade-dried for 7 days and then pulverized for Storage, protected from humidity.

2.2. Méthodes

2.2.1. Preparations of Extracts

The extracts were prepared according to methodologies used in the literature [31, 32].

Decoctates (5%) of the various solvents were obtained by extraction of leaves and roots three times under reflux for successive periods of 2h-1h30-1h. For each organ the 3 solutions are mixed.

Macerates (5%) are obtained after 24 hours. leaves and roots were macerated in different solvents (water – ethanol 70% – acetone 30%).

The extraction yield ρ (%) was expressed as follows:

$$\rho = (QA/QA0) * 100 \quad (1)$$

With QA: Amount of material extracted by the solvent
QA0: initial amount of plant material used

2.2.2. Phytochemical Screening

Phytochemical screening was carried out using the stepwise solvent method. The targeted compounds were alkaloids (with Bouchardat and Dragendorff methods), polyphenols (with $FeCl_3$ test), flavonoids (with cyanidin reaction), tannins (with Stiasny reaction followed by that of ferric chloride), saponins (by index of moss), cardiac glycosides (with Keller-kiliani methods), quinones (with Borntraeger reaction), sterols (with Liebermann-Buchard reaction) [33].

2.2.3. Inhibition of Protein déNaturation Using Egg Albumin Method

The anti-inflammatory potential of the aqueous, alcoholic and acetonic extracts of *Imperata cylindrica* was determined by the inhibition of the egg albumin denaturation method. The ability of *I. cylindrica* extracts to inhibit protein denaturation was determined by methods described in the literature with some modifications [34-36].

The reaction mixture consisted of 0.2 mL of egg albumin (1%), 4.78 mL PBS (pH 6.4) and 0.02 mL of diclofenac extract (1-2, 5-5-10 mg/mL). It was then incubated at $37^\circ C \pm 2$ for 15 min and then heated to $70^\circ C$ for 5 min. After cooling, the reading was taken with a spectrophotometer at 660nm against the blank consisting of PBS (pH 6.4). Distilled water and diclofenac served as negative and positive controls, respectively. The percentage of denaturation inhibition was calculated according to equation (2). The results of the extracts were compared to those of diclofenac.

$$\% \text{Anti-denat} = [100 * (\text{Abs extracts} / \text{Abs Contrôle}) - 1] \quad (2)$$

2.2.4. Inhibition of Lipid Peroxidation: TBARS Method

The reaction mixture was prepared with 10 μ L of extracts of *I. cylindrica* (10 mg/mL), 50 μ L of egg yolk solution (10% v/v in 1.15% KCl), 150 μ L of 20% acetic acid solution (pH 3.5) and 150 μ L of a solution of thiobarbituric acid (0.8% in 1.1% SDS). The volume of the reaction mixture is then adjusted to 400 μ L with distilled water. The whole is mixed

for 5 seconds by vortex then incubated at 95°C for 1 hour. At the end of the incubation time, 1mL of butanol is added. The mixture is stirred for 5 s and centrifuged at 1500 g for 5 min. The butanolic fraction is extracted and its absorbance is read at 532nm. Distilled water and ascorbic acid are used as negative (distilled water) and positive (0.2 mg/mL ascorbic acid) controls, respectively. The percentage of inhibition is calculated according to the following relationship [37].

$$\%Inh = 100*[1-(Abs\ contr\hat{o}le/ Abs\ extract)] \quad (3)$$

2.3. Statistical Analyses

Data are presented as mean \pm standard deviation (SD), and experiments were performed three times. The software (Microsoft Excel 2013) was used for measurements (means and standard deviations). Statistica 7.1 software was used for

analysis of variance (ANOVA). Comparison of the means was obtained by the LSD test (significance threshold = 0.05). Determination of IC₅₀ was made using the Graphpad software.

3. Results

Table 1. Extraction yield (%).

	Solvent	Leaves	Roots
Maceration	Water	3,86 \pm 0,35 ^h	6,89 \pm 0,26 ^f
	Alcohol	4,95 \pm 0,30 ^g	14,10 \pm 0,50 ^c
	Acetone	5,43 \pm 0,27 ^g	8,83 \pm 0,57 ^e
Decoction	Water	8,68 \pm 0,59 ^e	12,67 \pm 0,60 ^d
	Alcohol	7,91 \pm 0,28 ^e	18,33 \pm 0,76 ^a
	Acetone	8,58 \pm 0,36 ^e	15,27 \pm 0,52 ^b

On each column and each row, the yield values assigned different alphabetical letters are statistically different at $p < 0,05$ (ANOVA, LSD).

Table 2. Phytochemical profile of the different extracts of *Imperata cylindrica*.

	Roots			Leaves		
	Ethanol 70%	Eau	Acétone 30%	Ethanol 70%	Eau	Acétone 30%
Alkaloids	+	+	+	-	+	-
Polyphenols	+	+	+	+	+	+
Flavonoids	+	+	+	+	+	+
Polyterpenoids	-	-	-	+	+	+
Galic Tannins	+	+	+	+	+	+
Catechical Tannins	-	-	-	-	-	-
Saponins	-	+	+	-	+	+
Quinones	-	-	-	-	-	-

+: Presence, - Absence

Table 3. IC₅₀ of *Imperata cylindrica* concentration-dependent extracts.

EXTRAITS	IC ₅₀ mg/mL
FMO	5,76
FMA	13,64
FDA	8,94
FDC	6,48
RMO	12,47
RMC	11,08
RDO	5,94
Diclo	3,63


On each column and each row, the yield values assigned different alphabetical letters are statistically different at $p < 0,05$ (ANOVA, LSD). Diclo: diclofenac F: leaves; M: macération; D: decoction; R: roots

Table 4. Percentage of lipid peroxidation inhibition of *Imperata cylindrica* different extracts at 10 mg/mL and ascorbic acid at 0.2 mg/mL using egg yolk homogenate as medium.

EXTRAITS	% Inh
FMO	32,96 ^a \pm 0,65
FMA	32,43 ^a \pm 5,06
FMC	51,84 ^c \pm 2,85
FDO	39,33 ^{ab} \pm 0,39
FDA	69,81 ^{de} \pm 3,21
FDC	51,76 ^c \pm 2,26
RDO	76,33 ^{ef} \pm 6,14


EXTRAITS	% Inh
RDA	76,55 ^{ef} \pm 6,73
RDC	74,01 ^{ef} \pm 8,80
RMO	62,77 ^d \pm 5,27
RMA	43,52 ^b \pm 2,47
RMC	81,05 ^f \pm 3,16
AA (0,2mg/mL)	41,72 ^b \pm 4,63

The values expressed are the averages of the inhibition percentages, $n = 3$ replications, significant difference between the extracts at $p < 0,001$; Diclo: diclofenac F: leaves; M: macération; D: decoction; R: roots



Diclo: diclofenac F: leaves; M: macération; D: decoction; R: roots

Figure 1. Inhibition of protein (albumin) denaturation by different concentrations of various *Imperata cylindrica* roots extracts.



Diclo: diclofenac F: leaves; M: macération; D: decoction; R: roots

Figure 2. Inhibition of protein (albumin) denaturation by different concentrations of various *Imperata cylindrica* leaves extracts.

4. Discussion

The work explored extracts of *Imperata cylindrica* leaves and roots obtained from two methods (decoction and maceration) using three solvents (distilled water, ethanol 70%, acetone 30%) (Table 1). The extraction rate of leaves generally doubled with increasing temperature from 3.9% to 8% when moving from maceration to decoction. Solvent and temperature impacted extraction of the roots both in maceration and in decoction (Table 1).

The highest values are obtained by 70% ethanol ($14.10 \pm 0.50\%$ and $18.33 \pm 0.76\%$) followed by acetone ($8.83 \pm 0.57\%$ and $15.27 \pm 0.52\%$) and finally distilled water ($6.89 \pm 0.26\%$ and $12.67 \pm 0.60\%$). The increase in temperature probably impacted the solubility and diffusivity of the solvent and the breaking of bonds in plant cells for better extraction of molecules as mentioned in other works [38, 39]. Furthermore, it has been admitted that under the same extraction conditions the solvent is one of the most important factors that can impact the extraction yield [40].

Phytochemical screening of crude aqueous, ethanolic and acetoneic extracts of roots and leaves highlighted presence of several secondary metabolites (alkaloids, saponins, polyphenolic compounds) which are known to possess numerous medicinal properties in other studies [41-44]. Alkaloids are thought to have anti-inflammatory, antioxidant, antimicrobial and neuro-stimulating activities [45, 46]. Saponins are said to have anti-inflammatory and anti-diarrheal properties [47-49]. Other works have also shown preventive and/or therapeutic effects of polyphenols on metabolic diseases, neurodegenerative diseases, cell proliferation, inflammatory conditions and many other diseases [50, 51].

It is accepted that composition in secondary metabolites of medicinal plants is influenced by parameters as environment, period of harvest of the material, type of soil. Likewise extraction conditions (part of the plant material, particle size), extraction method, type and concentration of extraction solvents, solvent-plant ratio, extraction temperature and time, affect the rate and the quality of the extracted metabolites and

therefore their biological properties [22].

In vitro anti-inflammatory activity of *I. cylindrica* extracts was evaluated through the protective activity of the extract against albumin denaturation. It has been noted that this mode of action, in this case the inhibition of the denaturation of bovine serum albumin at pH 6.2-6.4, is one of the mechanisms followed by several non-steroidal anti-inflammatory drugs [10]. The higher the degree of inhibition of albumin denaturation, the greater its anti-inflammatory potential [11, 52-54]. In this test, the egg albumin was denatured by heat; the inhibitory effect of the different extracts is presented in Figure 1 and Figure 2.

The percentage of inhibition varies from 0% to 55.16% for the extracts and from 10.01% to 119.30% for Diclofenac®. An extract is considered to have anti-inflammatory power when it produces a percentage of inhibition > 20% [54]. The extracts in the study could be classified into three classes: extracts without effect (RDC), extracts exerting their maximum at low concentration (FDO, RDA, RMA, FMC) and extracts having a concentration-dependent anti-inflammatory power at 1mg/mL.

The RDA, RMA, FMC extracts with an inhibition percentage between (21.96% and 33.56%) would exert maximum anti-inflammatory activity. Indeed, work had noted that algae extracts (*Undaria pinnatifida*, *Fucus vesiculosus*, *Macrocystis pyrifera*, *Ascophyllum nodosum*, and *Laminaria japonica*) rich in fucoidanes reduced levels of pro-inflammatory markers in vitro both in PBMC and in THP-1 cells with maximum activity at low concentrations [55]. This effect has been attributed to the low molecular weight of the extracts, better absorption across the cell surface as well as a rapid mode of action. Most plant extracts, however, have anti-inflammatory activity concentration-dependent. In this case, determining the IC₅₀ makes it possible to classify them. The smaller the IC₅₀, the greater its inhibitory potential. On the basis of the IC₅₀ it was thus possible to estimate that the aqueous root decoction (RDO) and aqueous leaf macerate (FMO) with respectively 5.94 mg/mL and 5.76 mg/mL were the most effective extracts active in inhibiting denaturation of egg albumin. Their potential is, however, lower than that of our reference substance (Diclofenac) (3.63 mg/mL). The anti-inflammatory potential could be attributed to a phenolic derivative salicin structurally similar to Aspirin® and already isolated from flowers [56]. These results could justify the use of extracts of *I. cylindrica* against feverish conditions and malaria in a traditional environment [57].

During the work, the lipid supplier was egg yolk solution which contains between 31% and 35% lipids, mainly made up of linoleic and linolenic acids. The percentage of inhibition of the extracts is greater than 30% with a maximum activity ($81.05\% \pm 3.16$) at the level of the acetone macerate of the roots (RMC) which is however not statistically different from the values of the aqueous decoctions. (76.33%), alcoholic (76.55%) and acetone (74.33%) of the roots. The study also revealed that extracts from the root have a more interesting activity ($43.52b \pm 2.47$

to 81.05 ± 3.16 % than leaf extracts (32.43 ± 5.06 to 69.81 ± 3.21 %). It should be noted that most extracts have greater activity than ascorbic acid at 0.2 mg/mL. The results of the study suggest that *Imperata cylindrica* extracts could have a cytoprotective property by inhibiting or reducing lipoperoxidation of non-enzymatic origin in a lipid medium. Polysaccharides isolated from *I. cylindrica* have been shown to be effective in protecting against kidney damage by inhibiting lipid peroxidation, increasing antioxidant defenses and inhibiting the secretion of pro-inflammatory factors, regulating purine metabolism, PI3K-Akt, NF signaling pathways - κ B, mTOR and MAPK [58].

5. Conclusion

The present study on the anti-inflammatory and antioxidant activity of extracts of leaves and roots of *I. cylindrica* using different solvents and two extraction methods noted that the plant is rich in bioactive compounds. The study also identified alcohol as the best solvent for extracting these bioactive compounds. Furthermore, the study found that water was the suitable solvent to investigate the antilipoperoxidation and antidenaturation activities of proteins.

The optimal extraction parameter of antioxidant and anti-inflammatory compounds was: roots / water / decoction.

Author Contributions

'Conceptualization: Anne Marie Leticia Konan, Esmel Essis Lohoues, Methodology: Anne Marie Leticia Konan, Esmel Essis Lohoues, Félix Adjé, Amissa Augustin Adima, Validation: Anne Marie Leticia Konan, Esmel Essis Lohoues, Adou Kra Matthieu Kra, Writing - Original Draft Preparation: Anne Marie Leticia Konan, Félix Adjé, Koffi, Julien Golly, Kessé Phyllipe N'da, Writing - Review & Editing: Anne Marie Leticia Konan, Supervision: Esmel Essis Lohoues, Adou Kra Matthieu Kra, Amissa Augustin Adima.

ORCID

0000-0003-4613-295X (Anne Marie Leticia Konan)

0000-0002-4538-5747 (Félix Adjé)

0000-0001-5818-8374 (Amissa Augustin Adima)

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] Antonelli M and Kushner I, It's time to redefine inflammation. *FASEB Journal* 2017; 31: 1781- 1791.
- [2] Kristina H, Mrowka R, Inflammation -Dysregulated inflammatory response and strategies for treatment. *Acta Physiol* 2019; 226(3): e13284 <https://doi.org/10.1111/apha.13284>
- [3] Negishi Y, Shima Y, Takeshita T, Morita R, Harmful and beneficial effects of inflammatory response on reproduction: sterile and pathogen associated inflammation. *Immunol Med*; 2021. 44(2): 99 -115. <https://doi.org/10.1080/25785826.2020.1809951>
- [4] Shailasree S, Sampathkumara KK, Niranjana SR, Prakash HS, Bioactive Potential of Medicinal Plants from Western Ghats Region, India, *J. Herbs Spices Med. Plants*, 2014, 20: 3, 221-234.
- [5] Liu M, Kalbasi A, Beatty GL., *Functio Laesa: Cancer Inflammation and Therapeutic Resistance. J. Oncol. Pract*, 2017, 13 (3): 173-180. DOI: 10.1200/JOP.2016.020347.
- [6] Rossi J-F, Lu ZY, Massart C, Levon K, *Dynamic Immune/ Inflammation Precision Medicine: The Good and the Bad Inflammation in Infection and Cancer. Front. Immunol*, 2021, 12: 595722. doi: 10.3389/fimmu.2021.595722.
- [7] Kar Saptarshi, Kavdia Mahendra, (2012) Local Oxidative and Nitrosative Stress Increases in the Microcirculation during Leukocytes-Endothelial Cell Interactions. *PLoS ONE* 7(6): e38912. doi: 10.1371/journal.pone.0038912.
- [8] Ruiz-Ruiz JC, Matus-Basto AJ Acereto-Escoffé P, Segura-Campos MR, Antioxidant and anti-inflammatory activities of phenolic compounds isolated from *Melipona beecheii* honey, *Food Agric Immunol*, 2017, 28: 6, 1424-1437, DOI: 10.1080/09540105.2017.1347148.
- [9] Freigang S, The regulation of inflammation by oxidized phospholipids. *Eur. J. Immunol*, 2016, 46: 1818–1825. DOI: 10.1002/eji.201545676.
- [10] Di Gioia M, Zanoni I, Dooming Phagocyte Responses: Inflammatory Effects of Endogenous Oxidized Phospholipids. *Front Endocrinol*. 2021 12: 626842. doi: 10.3389/fendo.2021.626842.
- [11] Alamgeer UHH, Evaluation of in vitro and in vivo therapeutic efficacy of *Ribes alpestre* Decne in Rheumatoid arthritis. *Braz J Pharm Sci*; 2019, 55: e17832.
- [12] Di Meo S, Reed TT, Venditti P, Victor VM, Role of ROS and RNS Sources in Physiological and Pathological Conditions, *Oxid Med Cell Longev*, 2016, Article ID 1245049, 44 pages.
- [13] Sharifi-Rad Mehdi, Kumar Nanjangud V. Anil, Zucca Paolo, Varoni Elena Maria, Dini Luciana, Panzarini Elisa, Jovana Rajkovic, Patrick Valere Tsouh Fokou, Elena Azzini8 Ilaria Peluso, Abhay Prakash Mishra, Manisha Nigam, Youssef El Rayess, Marc El Beyrouthy, Letizia Polito, Marcello Iriti, Natália Martins, Miquel Martorell, Anca Oana Docea, William N. Setzer, Daniela Calina, William C. Cho, Javad Sharifi-Rad, 2020. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front. Physiol*; Volume 11 <https://doi.org/10.3389/fphys.2020.00694>
- [14] Martemucci Giovanni, Costagliola Ciro, Mariano Michele, D'andrea Luca, Pasquale Napolitano and Angela Gabriella D'Alessandro, 2022. Free Radical Properties, Source and Targets, Antioxidant Consumption and Health. *OxygeN*; 2/ 48–78. <https://doi.org/10.3390/oxygen2020006>
- [15] Taso OV, Philippou A, Moustogiannis A, Zevolis E, Koutsilieris M, Lipid peroxidation products and their role in neurodegenerative diseases. *Ann Res H*, 2019, doi: 10.21037/arh.2018.12.02.

- [16] Petrovic S, Arsic A, Ristic-Medic D, Lipid Peroxidation and Antioxidant Supplementation in Neurodegenerative Diseases: A Review of Human Studies. *Antioxidants*, 2020.; 9(11): 1128; <https://doi.org/10.3390/antiox9111128>
- [17] Sabanna PK, Ratan WR, Lipid Peroxidation: A Signaling Mechanism in Diagnosis of Diseases. *Accenting Lipid Peroxidation*. IntechOpen, 2021 Available at: <http://dx.doi.org/10.5772/intechopen.99706>.
- [18] Ramana KV, Srivastava S, Sharad SS, Lipid Peroxidation Products in Human Health and Disease *Oxid Med Cell Longev*, 2019, Article ID 7147235. <https://doi.org/10.1155/2019/7147235>
- [19] Dziabowska-Grabias K, Sztanke M, Zajac P, et al., Antioxidant Therapy in Inflammatory Bowel Diseases. *Antioxidants* 2021; 10, 412. <https://doi.org/10.3390/antiox10030412>:
- [20] Khanna RD, Karki K, Pande D, Negi R, Khanna RS, Inflammation, Free Radical Damage, Oxidative Stress and Cancer. *Microinflamm*, 2014, 1: 109. doi: 10.4172/2381-8727.1000109.
- [21] Michael Wink, 2022. Current Understanding of Modes of Action of Multicomponent Bioactive Phytochemicals: Potential for Nutraceuticals and Antimicrobials. *Annu. Rev. Food Sci. Technol.* 2022. 13: 337–59.
- [22] Zhang Q-W, Lin L-GG, Ye W-C, Techniques for extraction and isolation of natural products: a comprehensive review. *Chin Med*, 2018., 13,20. doi.org/10.1186/s13020-018-0177-x.
- [23] Brewer J. Stephen, 2008. Declines in plant species richness and endemic plant species in longleaf pine savannas invaded by *Imperata cylindrica*. *Biological Invasions*, 10: 1257-1264.
- [24] Kouakou JK, Yapi A, Alui AK, Akotto FO, Yao-Kouame A, Pédopaysage et distribution de *Imperata cylindrica* (L.) P. Beauv. (Poaceae) dans deux Agro-écosystèmes de la Côte d'Ivoire: Abidjan et Bouaké. *Intern J. Innov. Sc. Res.* 2016; 22(1): 238-249.
- [25] Koné WM, Koffi AG, Bomisso EL, Tra Bi FH, Ethnomedical Study And Iron Content Of Some Medicinal Herbs Used In Traditional Medicine In Côte d'Ivoire For The Treatment Of Anaemia. *Afr J Tradit Complement Altern Med.*, 2012, 9(1): 81-87.
- [26] Parvathy NG, Padma R, Renjith V, Rahate KP, Saranya TS, Phytochemical screening and anthelmintic activity of methanolic extract of *imperata cylindrica*. *Int J Pharm Pharm Sci*, 2012. 4: 232-234.
- [27] Ruslin, Asmawi ZM, Rianse U et al., Antihypertensive activity pf Alang Alang (*Imperata cylindrica* (L.) Beauv. Root Methanolic extract on male wista rat. *Int. J. Res. Pharm. Sci.*, 2013, 4(4): 537-542.
- [28] Liu R-h, Chen S-S, Ren G, Shao F, Huang H-l, Compounds from Roots of *Imperata cylindrica* var. major. *Chin Herb Med*, 2013, 5: 240-243.
- [29] An H-J, Nugroho A, Song B-Mi, Park H-J, Isoeugenin, a Novel Nitric Oxide Synthase Inhibitor Isolated from the Rhizomes of *Imperata cylindrica*. *Molec*, 2015, 20: 21336–21345.
- [30] Xuan L, Bin-Feng Z, Li Y, Gui-Xin C, Zheng-Tao W, Four new compounds from *Imperata cylindrica*. *J Nat Med*, 2014, 68: 295–301.
- [31] Zhou X-r, Wang J-H, Jiang B, Shang J, Zhao C-Q, A Study of extraction process and in vitro antioxydant activity of total phénols from *Rhizoma imperatae*. *Afr J Tradit Complement Altern Med*, 2013,; 10(4): 175-178. doi.org/10.4314/ajtcam.v10i4.28.
- [32] Abe E, Delyle SG, Alvarez JC, Extraction liquide-liquide: théorie, applications, difficultés / Liquid-liquid extraction: theory, applications and difficulties. *Ann Toxicol Anal*, 2010, 22(2): 51-59.
- [33] Nancy RIM, Meenashree B, Vasanthi VJ, Screening of Antibacterial Activity and Qualitative and Quantitative analysis of Phytochemicals in *Vitex trifolia*, *Int. J. Curr. Microbiol. App. Sci.*, 2014, 3: 425-431.
- [34] Gunathilake KDPP, Ranaweera KKDS, Vasantha Rupasinghe HP, In Vitro Anti-Inflammatory Properties of Selected Green Leafy Vegetables. *Biomedicines*, 2018. (6): 107. doi: 10.3390/biomedicines6040107.
- [35] Banerjee S, Chanda A, Adhikari A, Das A, Biswas S, Evaluation of Phytochemical Screening and Anti Inflammatory Activity of Leaves and Stem of *Mikania scandens* (L.) Wild. *Ann Med Health Sci Res*, 2014. 4(4): 532–536.
- [36] Ullah HMA, Zaman S, Juhara F, et al., Evaluation of antinociceptive, in-vivo & in-vitro anti-inflammatory activity of ethanolic extract of *Curcuma zedoaria* rhizome. *BMC Complem Altern Med*, 2014. 14: 346. <http://doi.org/10.1186/1472-6882-14-346>.
- [37] Dissanayake DP, Abeytunga DTU, Vasudewa NS, Ratnasooriya WD. Inhibition of lipid peroxidation by extracts of *Pleurotus ostreatus*. *Pharmacogn Mag*, 2009, 4(19): 266-271.
- [38] Chaves JO, de Souza MC, da Silva LC, et al., Extraction of Flavonoids From Natural Sources Using Modern Techniques. *Front. Chem*, 2020, 8: 507887. doi: 10.3389/fchem.2020.507887.
- [39] Lezoul NEH, Belkadi M, Habibi F, Guillén F, Extraction Processes with Several Solvents on Total Bioactive Compounds in Different Organs of Three Medicinal Plants. *Molec*, 2020. 25(20): 4672. doi: 10.3390/molecules25204672.
- [40] Truong D-H, Nguyen DH, Ta NTA, et al., Evaluation of the Use of Different Solvents for Phytochemical Constituents, Antioxidants, and In Vitro Anti-Inflammatory Activities of *Severinia buxifolia*. *J food qual*, 2019 Article ID 8178294 | <https://doi.org/10.1155/2019/8178294>.
- [41] Iram F, Sobia K, Tariq M, Evaluation of Biological Potential of Selected Species of Family Poaceae from Bahawalpur, Pakistan. *BMC Compl Altern. Med.*, 2018. 18, Article No. 27. <https://doi.org/10.1186/s12906-018-2092-1>.
- [42] Lalthanpuui PB, Zarzokimi, Lalchhandama K, *Imperata cylindrica*: A Noxious Weed of Pharmacological Potentials In Advances in Engineering Research Perspective and Trends in the Development of Science Education and Research. *Atlantis Press*, 2018, Paris, 173-177. <https://doi.org/10.2991/msc-18.2018.28>.
- [43] Sudha R, Kaleena PK, Babu M, Janaki A, Velu K, Elumalai D, Phytochemical Screening, Antioxidant and Anticancer Potential of *Imperata cylindrica* (L.) Raeusch against Human Breast Cancer Cell Line (mcf-7). *Int. j. pharm. biol. Sci*, 2018, 8: 938-945.

- [44] Bashige VC, Bakari AS, Okusa PN, Kalonda EM, Lumbu JBS, Criblage phytochimique et activité antimicrobienne de six rhizomes comestibles utilisés en médecine traditionnelle à Lubumbashi (RDC). *Int. j. biol. chem. Sci.*, 2020 14: 1367-1380. <https://doi.org/10.4314/ijbcs.v14i4.16>
- [45] Hussain G, Rasul A, Anwar H, et al., Role of Plant Derived Alkaloids and Their Mechanism in Neurodegenerative Disorders. *Int j pharm boil sci*, 2018. 14: 341-357. <https://doi.org/10.7150/ijbs.23247>.
- [46] Sicong L, Xin L, Xiaoran C, Lei B, Research Progress on Anti-Inflammatory Effects and Mechanisms of Alkaloids from Chinese Medical Herbs. *Evidence-Based Complement Altern Med*, Volume 2020, Article ID 1303524, 10 pages <https://doi.org/10.1155/2020/1303524>
- [47] Johnson AM, 2013. Saponins as Agents Preventing Infection Caused by Common Waterborne Pathogens. Dissertation, University of Texas, Austin, 126 p.
- [48] Shi Y, Guo R, Wang X et al., The Regulation of Alfalfa Saponin Extract on Key Genes Involved in Hepatic Cholesterol Metabolism in Hyperlipidemic Rats. *Plos One*, 2014., 9: e88282. <https://doi.org/10.1371/journal.pone.0088282>.
- [49] Sun X-B, Liu Y-P, Yang Y-Y, Liu X-Y, Xiang D-X, Anti-Arthritic Effect of Total Saponins from *Clematis henryi* Oliv. on Collagen-Induced Arthritis Rats. *Eur J Inflamm*, 2016, 14: 71-77. <https://doi.org/10.1177/1721727X16644448>.
- [50] Dover A, Patel N, Park KT, Rapid Cessation of Acute Diarrhea Using a Novel Solution of Bioactive Polyphenols: A Randomized Trial in Nicaraguan Children. *Peer J* 2015, 3: e969. <https://doi.org/10.7717/peerj.969>.
- [51] Cory H, Passarelli S, Szeto J, Tamez M, Mattei J, The Role of Polyphenols in Human Health and Food Systems: A Mini-Review. *Front Nutr* 2018, 5, article 87. <https://doi.org/10.3389/fnut.2018.00087>.
- [52] Yani DF, Fatahillah R, Inflammatory activity os ethanol extract and ethyl acetate fraction of kebuil (*Caesalpinia bonduc* L.) seed coat against inhibition of proteindenaturation. *J Kimia Riset*, 2022, 7(1): 2022 1-81.
- [53] Cano-Avendaño BA, Carmona-Hernandez JC, Rodriguez RE, Taborda-Ocampo G, González-Correa CH, Chemical Properties of Polyphenols: A Review Focused on Anti-Inflammatory and Anti-Viral Medical Application. *Biomed*, 2021, 41: 3-8. <https://doi.org/10.51248/v4i1i.524>.
- [54] Heendeniya SN, Ratnasooriya DW, Pathirana RN., In vitro investigation of anti-inflammatory activity and evaluation of phytochemical profile of *Syzygium caryophyllatum*, *j Pharmacogn Phytochem*, 2018; 7(1): 1759-1763.
- [55] Ahmad T, Eapen MS, Ishaq M, et al., Anti-Inflammatory Activity of Fucoidan Extracts In Vitro. *Mar. Drugs*, 2021, 19: 702. [doi.org/ 10.3390/md19120702](https://doi.org/10.3390/md19120702).
- [56] Abdullah FO, Hussain FHS, Sardar AS, Vita-Finzi P, Vidari G, Phytochemistry and Ethnopharmacology of Medicinal Plants Used on Safeen Mountain in the Kurdistan Region of Iraq, *Natural Product Communications*, 2016, 11(12): 1923-1927.
- [57] Lepetu, JP, *Imperata cylindrica* (L.) Raeusch. [Accessed online 16 October 2023] Record from PROTA4U. Brink, M. & Achigan-Dako, E. G. (Editors). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), 2011, Wageningen, Netherlands. <http://www.prota4u.org/search.asp>.
- [58] Guichardant, Ping C, Miao L, et al., Functional lipidomics of oxidized products from polyunsaturated fatty acids.. *Chem Phys Lipids*, 2011, 2164(6): 544-548.