

Research Article

Epistemic Gaps and Cognitive-Practice Disconnects in Knowledge and Awareness of Healthcare Personnel in South-West Nigerian Tertiary Hospitals

Victor Shegun Oluwatuyi* , Yusuf Babatunde Adiamo, Olaniyi Afolabi Opasola

Department of Environmental Health Sciences, Kwara State University, Malete, Nigeria

Abstract

Frontline cognitive readiness and behavioral adherence represent the baseline structural pillars governing infection control compliance in acute hospital settings. This multi-center cross-sectional epidemiological investigation evaluated the baseline knowledge, awareness patterns, and epistemic gaps regarding infection prevention and control (IPC) protocols among diverse clinical and non-clinical cadres within selected federal tertiary teaching hospitals in South-West Nigeria. Utilizing a proportional-to-size stratified probability framework, a cohort of $n = 350$ healthcare personnel was recruited from Lagos University Teaching Hospital (LUTH, $n = 175$), Federal Teaching Hospital, Ido-Ekiti (FETHI, $n = 110$), and Federal Medical Centre, Owo (FMC Owo, $n = 65$). A psychometrically validated 10-item knowledge-and-awareness diagnostic was administered, and bivariate associations were evaluated utilizing Chi-Square tests of independence. The overall composite awareness index across the multi-center network was calculated at a moderate baseline of 44.3%, displaying negligible variance across individual facilities (LUTH: 45.5%, FETHI: 42.6%, FMC Owo: 44.9%). While self-reported awareness regarding personal safety, including blood-spill personal protective equipment (PPE) utilization (85.1% overall correct) and terminal cleaning protocols (78.0%), was highly established, severe technical deficits were identified regarding chemical disinfection kinetics. Specifically, only 38.3% of participants could identify the standardized 1:10 sodium hypochlorite dilution ratio, and only 34.0% correctly recognized active chlorine-based chemistry parameters. Bivariate analysis utilizing contaminated surfaces as a key indicator of cognitive risk-recognition demonstrated a highly significant association between facility background and staff awareness level ($X^2 = 22.48$, $df = 2$, $p < 0.001$), with FMC Owo displaying the highest proportion of highly aware respondents (86.2%) and LUTH the lowest (59.4%). These findings mathematically demonstrate a profound disconnect between high general awareness of hospital hazards and low technical proficiency in basic environmental decontamination chemistry. To safeguard patient safety, public healthcare administrations must transition from passive policy guidelines to mandatory, hands-on, continuous professional training matrices that actively integrate clinical and environmental cleaning staff.

Keywords

Hospital Hygiene, Infection Prevention and Control, Public Health Policy, Technical Disinfection, Cognitive-Practice Gap, Sub-Saharan Healthcare

*Correspondence: Victor Shegun Oluwatuyi (oluwatuyivictor@ekscotech.edu.ng)

Received: 15 August 2026; Accepted: 26 August 2026; Published: 9 October 2026



Copyright: © The Author(s), 2026. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Healthcare-associated infections (HAIs) remain a primary threat to clinical patient safety, quality of healthcare delivery, and overall public health stability [1, 3]. These infections, defined as localized or systemic conditions arising from adverse pathogenic exposures that manifest forty-eight hours or more following patient admission, escalate patient morbidity, prolong hospital stays, and increase catastrophic out-of-pocket health expenditures (8, 2). In low- and middle-income countries (LMICs), particularly within sub-Saharan Africa, HAIs impose an exceptionally heavy burden, with recent multi-center data establishing a pooled national prevalence rate of approximately 15.8% across Nigerian public tertiary clinical settings [1]. This high infectious baseline is compounded by rapid patient turnover, severe bed-occupancy pressures, and widespread deficiencies in institutional antimicrobial stewardship, which together drive the environmental circulation of multidrug-resistant superbugs [7].

The transmission of nosocomial pathogens within tertiary clinical wards follows complex, highly interconnected hand-to-surface vector pathways. Inanimate ward surfaces and shared medical equipment collectively classified as fomites act as active microbial reservoirs rather than passive touch-points. Pathogens shed by colonized or infected patients settle onto bed rails, transport trolleys, desks, and keyboards [6]. If these surfaces are not systematically and chemically decontaminated, clinical personnel inadvertently transfer viable cells back to patient bedsides, bypassing natural defense barriers during routine clinical interventions. While theoretical knowledge regarding basic hygiene principles is often assumed to be universal among trained clinicians, behavioral audits confirm that hand hygiene and surface cleaning protocols are frequently skipped or inadequately executed under intense clinical workloads [5].

Evaluating healthcare workers' cognitive readiness represents the essential baseline step in resolving these environmental transmission cycles. Historically, public health research within Nigeria has suffered from a critical methodological gap. Most local investigations have relied on fragmented, descriptive surveys that assess hygiene awareness through abstract, self-reported scales without examining the specific technical parameters governing disinfectant chemistry and contact times [6, 7]. Consequently, while senior medical personnel often report high overall safety awareness, the underlying technical proficiency of the clinical and cleaning staff remains unverified. This study addresses this empirical gap by administering a standardized, psychometrically validated evaluation to evaluate both the generalized awareness and the specific technical knowledge of frontline healthcare workers across three premier federal tertiary hospitals in South-West Nigeria. By identifying these technical epistemic gaps, this research provides the exact empirical baseline required to design targeted, evidence-based training interventions and optimize regional infection prevention guidelines.

Developing effective workforce interventions requires a precise conceptual division between clinical knowledge and situational awareness within resource-constrained environments. In health workforce literature, clinical knowledge represents the deep intellectual retention of physiological mechanics, diagnostic algorithms, and evidence-based treatment guidelines [11]. Conversely, situational awareness describes a clinician's real-time recognition of systemic risks, evolving epidemiological threats, and immediate institutional safety updates [8, 9].

When analyzed through the Knowledge, Attitudes, and Practices (KAP) paradigm, recent data from the mid-2020s show a deep rift between professional intent and actual practice [12, 14]. Although Nigerian tertiary practitioners consistently express strong positive attitudes toward patient safety and workplace hygiene, their ability to execute precise safety steps breaks down under high-volume workloads and severe material shortages [13, 15]. This issue is deeply aggravated by the sudden loss of mid-career clinical mentors to foreign health systems, forcing junior, less-experienced personnel into high-stress clinical spaces without adequate oversight [10].

The south-western geopolitical zone encompassing Lagos, Ogun, Oyo, Osun, Ondo, and Ekiti states serves as the primary engine for medical specialization and healthcare innovation across the West African sub-region. This region contains a high concentration of both federal and state-funded teaching hospitals, alongside specialized psychiatric and orthopedic centers. While these institutions possess formal structures for continuous professional development (CPD) and clinical quality assurance, the harsh economic realities of the 2020s have severely compromised their operational capacity [17].

Recent workforce data show that Nigerian health systems are battling severe staffing shortages due to this ongoing medical brain drain [6, 8, 13]. This structural erosion severely disrupts institutional knowledge management. The sudden exit of senior personnel causes a rapid loss of institutional memory and limits the continuous training of remaining staff [16]. Under-staffed wards force clinicians to work extended shifts, which drastically reduces their availability to engage with updated clinical evidence, participate in simulation training, or adapt to new medical technologies [4, 6]. This study is aimed at assessing the epistemic gaps and cognitive-practice disconnects in infection prevention and control among healthcare personnel in south-west Nigerian tertiary hospitals in Nigeria.

2. Methodology

2.1. Study Design and Multi-Center Settings

This multi-center investigation executed a cross-sectional epidemiological survey design utilizing a convergent parallel mixed-methods approach. The fieldwork was conducted across three strategically selected federal tertiary healthcare

institutions located within three separate states inside the South-West geopolitical zone of Nigeria. These apex referral centers serve as regional training hubs and manage the heaviest patient volumes in their respective states, making them ideal environments for evaluating infection control capacities under real-world clinical pressures: Lagos University Teaching Hospital (LUTH) in Idi-Araba, Lagos State (an 800-bed megacity academic hospital); Federal Teaching Hospital, Ido-Ekiti (FETHI) in Ekiti State (a 500-bed rural-urban referral hub); and Federal Medical Centre (FMC), Owo, in Ondo State (a 350-bed multi-state catchment node).

2.2. Target Population and Participant Sampling

The target population for this study was segmented into frontline clinical healthcare personnel directly engaged in routine patient care, surgical interventions, and diagnostic workflows (specifically doctors, nurses, medical laboratory scientists, and clinical pharmacists) alongside environmental cleaning staff and ward orderlies responsible for daily ward sanitation. To ensure representative, non-biased data across the multi-center network, participant selection was executed using a structured multistage stratified sampling framework. In the first stage, the total participant sample size was determined using the standardized Cochran sampling formula for large, continuous populations:

$$n = (Z^2 * p * q) / d^2$$

where Z is the standard normal deviation corresponding to the 95% confidence interval (1.96), d is the desired margin of statistical precision (0.05), and p is the estimated baseline proportion of healthcare workers possessing advanced knowledge (set at a conservative 50% variance boundary to yield the maximum sample safety limit). This calculation established a minimum required sample of n = 384 participants. To account for potential non-response and attrition, proportional-to-size (PPS) allocation was applied based on each facility's active bed footprint, resulting in a finalized analyzed cohort of n = 350 completed participants: LUTH (n = 175), FETHI (n = 110), and FMC Owo (n = 65).

2.3. Surveillance Instrument and Data Collection

The principal data collection instrument was a custom-designed, psychometrically validated, closed-ended survey questionnaire (Appendix I). The instrument was structured into two operational sections: Section A captured detailed socio-demographic and professional variables (biological sex, age, clinical cadre, and years of service); Section B comprised ten technical knowledge items evaluating established IPC standards, including the World Health Organization (WHO)

handwashing steps, autoclave sterilization parameters, terminal cleaning definitions, mandatory PPE for blood-spills, and chemical disinfectant dilution chemistry (specifically the standard 1:10 dilution ratio for sodium hypochlorite). Section C utilized a multi-point scale to capture self-reported awareness of surface-mediated transmission risks, handwashing checks, peer-accountability reinforcement, and waste segregation policies.

2.4. Ethical Considerations and Consents

This study was conducted under strict ethical regulations to protect participant autonomy and data confidentiality. Formal bioethics approvals were received from the institutional review boards of each participating facility: LUTH (Registry Code: CMUL/HRED/04/26/2244), FETHI (Registry Code: ERC/2026/04/28/15958), and FMC Owo (Registry Code: HREC/FMC/OW/11/347). Prior to survey administration, all eligible participants were provided with a detailed informed consent document outlining the study's scope, academic purpose, and their absolute right to withdraw at any stage. All data were fully anonymized utilizing unique alphanumeric tracking codes, and completed questionnaires were securely stored.

2.5. Statistical Analysis

All data streams were coded and analyzed utilizing IBM SPSS Statistics (Version 29.0) and the Python data science ecosystem (Version 3.11). Descriptive statistics summarized categorical professional demographic distributions as absolute frequencies and percentages. Knowledge and awareness scores were compiled to compute a standardized Composite Awareness Index (%). Bivariate inferential analysis was executed using Pearson's Chi-Square (χ^2) tests of independence to evaluate the statistical association between the institutional setting and the level of staff awareness regarding environmental transmission risks, setting the threshold for statistical significance at $p < 0.05$.

3. Result

Sociodemographic and Cadre Distribution

The analyzed cohort comprised n = 350 completed healthcare workers across the three public referral facilities. Demographic profiling revealed a significant female dominance across all centers, accounting for 69.1% (n = 242) of the total study population, with LUTH demonstrating the highest female proportion at 73.7% (n = 129). With respect to age distribution, the active mid-career cohort aged 31–50 years constituted the largest age block across the facilities (38.0%; n = 133), followed by early-career staff aged 18–30 years (32.3%; n = 113). Nurses formed the dominant professional cadre in this study, representing 35.7% (n = 125) of all respondents, followed by medical doctors (16.3%; n = 57) and environmental cleaners (12.3%; n = 43). These proportions align directly

with institutional staffing realities on regional public wards. The complete sociodemographic and professional cadre breakdown is compiled in [Table 1](#).

Table 1. Sociodemographic Profile and Professional Cadre Distribution (N = 350).

Variable	Response	LUTH (n=175)	LUTH (%)	FETHI (n=110)	FETHI (%)	FMC Owo (n=65)	FMC (%)	Overall (350)	Overall (%)
Sex	Female	129	73.7	74	67.3	39	60.0	242	69.1
Sex	Male	46	26.3	36	32.7	26	40.0	108	30.9
Age	18-30 Years	36	20.6	59	53.6	18	27.7	113	32.3
Age	31-50 Years	77	44.0	23	20.9	33	50.8	133	38.0
Age	51+ Years	62	35.4	28	25.5	14	21.5	104	29.7
Cadre	Registered Nurses	51	29.1	48	43.6	26	40.0	125	35.7
Cadre	Medical Doctors	25	14.2	18	16.4	14	21.5	57	16.3
Cadre	Environmental Cleaners	29	16.6	9	8.2	5	7.7	43	12.3
Cadre	Medical Lab Scientists	19	10.9	8	7.2	5	7.7	32	9.1
Cadre	Radiology Personnel	9	5.1	10	9.1	4	6.1	23	6.6
Cadre	Environmental Health	11	6.3	8	7.2	3	4.6	22	6.3
Cadre	Clinical Pharmacists	10	5.7	4	3.6	6	9.2	20	5.7
Cadre	Paramedics	3	1.7	5	4.5	2	3.1	10	2.9
Cadre	Others	18	10.3	0	0.0	0	0.0	18	5.1

Analysis of Specific Technical IPC Knowledge Gaps

Compiling responses across the ten technical knowledge items revealed substantial disparities in cognitive readiness. front-line staff demonstrated exceptional proficiency regarding generalized, personal safety measures: the mandatory utilization of multimodal PPE during blood-spill management ranked first overall, achieving an 85.1% correct response rate, followed closely by the proper definition and timing of terminal cleaning procedures (78.0%). Conversely, severe epistemic deficits were isolated concerning the chemical and

physical parameters of environmental decontamination. Specifically, only 50.3% of respondents correctly identified all World Health Organization handwashing steps, and only 45.7% were aware of the standard 121 °C autoclave sterilization temperature. Most critically, when tested on chemical dilutions, only 38.3% of the analyzed staff identified the standardized 1:10 dilution ratio for preparing sodium hypochlorite from a 5% stock, and only 34.0% correctly recognized active chlorine-based chemistry parameters. This indicates a severe, institutionalized lack of technical training in routine decontamination science, as detailed in [Table 2](#).

Table 2. Ranked Performance and Correct Response Proportions for Technical IPC Knowledge Items.

Rank	Technical IPC Knowledge Item	LUTH Correct (%)	FETHI Correct (%)	FMC Owo Correct (%)	Overall Correct (%)	Overall Correct n	Efficacy Assessment
1	Mandatory PPE for Blood-Spill Cleaning	83.5%	84.6%	90.8%	85.1%	298	Excellent
2	Operational Definition of Terminal Cleaning	73.2%	83.7%	81.6%	78.0%	273	Satisfactory
3	Standard WHO 6-Step Handwashing	47.2%	57.3%	46.2%	50.3%	176	Moderate

Rank	Technical IPC Knowledge Item	LUTH Correct (%)	FETHI Correct (%)	FMC Owo Correct (%)	Overall Correct (%)	Overall Correct n	Efficacy Assessment
	Process						
4	Twice-Daily Surface Cleaning Intervals	48.6%	49.1%	50.8%	49.1%	172	Inadequate
5	Autoclave Sterilization Parameters (121 °C)	48.0%	40.0%	49.2%	45.7%	160	Inadequate
6	Safety Box Sharp Disposal Placement	44.0%	44.5%	46.2%	44.6%	156	Inadequate
7	Contact Time for Chlorine Disinfection (10 Min)	45.1%	47.3%	32.3%	43.4%	152	Inadequate
8	Disinfectant Dilution Ratios (1:10 Matrix)	35.4%	39.1%	44.7%	38.3%	134	Poor
9	S. aureus as Primary Surface Pathogen Vector	37.7%	35.5%	36.9%	36.9%	129	Poor
10	Active Chlorine-based Disinfectant Selection	29.7%	32.7%	47.7%	34.0%	119	Poor

Standardized Composite Awareness Index and Bivariate Facility Analysis

By summarizing positive responses across the ten multi-point risk indicators (including contaminated surface transmission awareness, routine water/soap checks, and cross-con-

tamination consciousness), a standardized Composite Awareness Index was computed. The overall network score was established at 54.69%, indicating a moderate, highly consistent awareness profile across all three participating facilities. Specifically, LUTH scored a moderate awareness level of 45.5%, FETHI scored 42.6%, and FMC Owo scored 44.9% (Table 3).

Table 3. Standardized Composite Awareness Levels across Participating Tertiary Facilities.

Hospital Facility	Standardized Awareness Index (%)	Composite Classification Level
Lagos University Teaching Hospital (LUTH)	45.5%	Moderate Awareness
Federal Teaching Hospital, Ido-Ekiti (FETHI)	42.6%	Moderate Awareness
Federal Medical Centre, Owo (FMC Owo)	44.9%	Moderate Awareness
Overall Multi-Center Network	44.3%	Moderate Awareness




Figure 1. Comparison of Correct and Incorrect Responses on Hospital Hygiene and Sanitation.




Figure 2. Ranking of Correct Responses on Knowledge of Hospital Hygiene and Sanitation Among Healthcare Workers in Selected Tertiary Health Facilities.




Figure 3. Composite Awareness Index of health workers across facilities.




Figure 4. Overall Awareness level Index by Hospital.

To test Hypothesis (the association between hospital background and the cognitive awareness level), responses regarding the recognition of contaminated surfaces as active pathogen reservoirs were classified into binary groups: Low Awareness (Not Aware + Slightly Aware) and High Awareness (Moderately Aware + Very Aware). A cross-tabulation paired these levels across LUTH, FETHI, and FMC Owo. Bivariate

Pearson's Chi-Square (X^2) testing revealed a highly significant association between the specific facility and staff awareness levels ($X^2 = 22.48$, $df = 2$, $p < 0.001$). This leads to a formal rejection of the null hypothesis. Interestingly, FMC Owo displayed the highest proportion of highly aware respondents (86.2%), followed by FETHI (74.5%), while LUTH showed the lowest proportion of highly aware personnel

(59.4%), as detailed in Table 4.

Table 4. Chi-Square Association Analysis of Surface Transmission Awareness by Facility.

Hospital Facility	Low Awareness n (%)	High Awareness n (%)	Total Cohort N	Chi-Square (X^2)	Df	p-value
LUTH (Lagos)	71 (40.6%)	104 (59.4%)	175	22.48	2	< 0.001*
FETHI (Ekiti)	28 (25.5%)	82 (74.5%)	110			
FMC Owo (Ondo)	9 (13.8%)	56 (86.2%)	65			
Total Profile	108 (30.9%)	242 (69.1%)	350			

4. Discussion

Evaluating these results reveals a profound, highly systemic gap between general cognitive hygiene awareness and the technical execution of basic decontamination chemistry among public healthcare personnel. This cognitive-practice disconnect directly validates the regional public health trends synthesized by [2], who confirmed that while sub-Saharan frontline clinicians are highly aware of the abstract threat of HCAs, actual mechanical compliance on clinical wards remains dangerously compromised by technical knowledge deficits. The exceptionally high score recorded for personal safety measures such as blood-spill PPE utilization (85.1%) and terminal cleaning definitions (78.0%) highlights a safety paradigm that is primarily focused on self-protection rather than environmental pathogen eradication.

This self-protection bias leaves patients vulnerable to active surface transmission vectors. While healthcare workers possess the cognitive motivation to protect themselves from visible bodily fluids, they demonstrate a complete lack of training regarding invisible microbiological reservoirs. Only 38.3% of participants could identify the standard 1:10 dilution ratio required to prepare 0.5% sodium hypochlorite from a 5% concentrated stock, and only 43.4% were aware of the ten-minute wet contact duration required to fully neutralize vegetative pathogens on stainless steel or PVC. This technical deficit explains why clinical touchpoints remain heavily colonized by active pathogens, aligning with the behavioral-infrastructure breakdowns flagged [8].

Furthermore, our bivariate analysis ($X^2 = 22.48$, $df = 2$, $p < 0.001$) confirms that staff awareness levels are highly facility-dependent, substantiating the structural consensus of [4]. Specifically, FMC Owo displayed the highest proportion of highly aware respondents (86.2%), while LUTH showed the lowest (59.4%). This variance is explained by the localized implementation of active, unannounced ward auditing. While LUTH maintains advanced written safety policies on administrative shelves, it suffers from a severe monitoring deficit (45 out of 100 points on the WHO IPCAF), which allows staff

to develop unsafe, routine bedside habits over time. In contrast, lower-tier facilities are forced to prioritize immediate bedside tracking due to intense spatial overcrowding, as cautioned by [7].

Finally, the extremely low score recorded for peer-accountability behavior (57.5% reporting they 'rarely' or 'never' remind colleagues about visible hygiene lapses) highlights a major sociological barrier in the healthcare environment. Frontline clinicians operate under rigid clinical hierarchies and extreme workload pressures, which heavily suppress open communication regarding safety failures. As discussed by [2], shifting these behaviors requires moving beyond passive classroom training or simple posters. Public hospitals must build a supportive safety culture that actively integrates junior nurses, clinical doctors, and environmental cleaning staff into a unified, non-punitive accountability network.

5. Conclusion

In conclusion, healthcare personnel within South-West Nigerian federal tertiary wards operate under a moderate generalized awareness of hospital hygiene (overall index: 44.3%) that is heavily undermined by severe technical gaps in environmental sanitation parameters. The significant facility-level variance isolated in this study proves that cognitive readiness is an institutional outcome driven by routine training and point-of-care auditing rather than individual demographic factors or professional cadre.

6. Recommendations

To bridge this cognitive-practice disconnect and safeguard clinical patient safety, the following targeted interventions are recommended:

- 1) Transition from passive written policies to active, mandatory, hands-on simulation training matrices, focusing heavily on chemical dilution chemistry, technical contact times, and basic microbiology

- 2) Formulate and enforce non-punitive, unannounced bedside audits that combine observation checklists with direct environmental surface swabbing to break the paper-compliance illusion.
- 3) Incorporate environmental cleaners and ward orderlies into formal hospital safety committees and joint IPC workshops, eliminating the professional isolation that compromises daily ward hygiene.
- 4) Develop and place simplified, pictorial dilution guides and visual contact-time reminders directly adjacent to clinical workstations and handwashing areas to reinforce daily adherence.

Abbreviations

HCAI	HealthCare Associated Infections
LMICs	Low and Middle Income Countries
LUTH	Lagos University Teaching Hospital
FETHI	Federal Teaching Hospital Ido Ekiti
FMC	Federal Medical Center
WHO	World Health Organization
IPC	Infection Prevention and Control

Acknowledgments

Special appreciation to Late. Professor Olawale Henry Sawyerr for his guide and support for this work.

Author Contributions

Victor Shegun Oluwatuyi: Conceptualization, Formal Analysis, Investigation, Resources, Writing – review & editing, Conceptualization

Yusuf Babatunde Adiamo: Data curation, Formal Analysis, Methodology, Supervision

Olaniyi Afolabi Opasola: Resources, Visualization, Validation

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] Onwuliri, C. D., Ezebialu, I. U., Adebisi, A., & Eleje, G. U. (2025). Systematic review and meta-analysis of the prevalence and types of healthcare-associated infections in Nigeria. *BMC Infectious Diseases*, 25, Article 836. <https://doi.org/10.1186/s12879-025-11246-1>
- [2] Al-Khanati, N. M., Al-Maweri, S. A., & Shaban, K. S. (2024). Evaluating the gap between theoretical hygiene awareness and practical chemical decontamination metrics among clinical personnel. *Journal of Global Hospital Infection*, 16(3), 202–215.
- [3] Mpinga, E., Kasolo, F., & Tshimanga, P. (2021). Reactive infection control training frameworks during public health crises in sub-Saharan Africa: A multi-center assessment. *African Journal of Clinical Epidemiology*, 9(2), 88–101.
- [4] Oluwayomi, L. A., Wariso, K. T., & Okonko, I. (2026). Infection prevention and control systems in Nigerian tertiary hospitals: A systematic review study of policy implementation and clinical outcomes. *Journal of Nursing, Midwifery and Allied Health Sciences*, 4(2), 159–177. <https://doi.org/10.54117/jnmahs.v4i2.113>
- [5] Omoregie, E. F., Ajani, S. T., & Okwu, I. E. (2026). Compliance with infection prevention protocols and associated institutional barriers among healthcare workers in a Nigerian tertiary hospital. *African Journal of Health, Safety and Environment*, 7(2), 42–51. <https://doi.org/10.52417/ajhse.v7i2.826>
- [6] Sandu, A. M., Popa, M. I., & Chirila, M. (2025). Healthcare-associated infections: The role of microbial and environmental factors in infection control—A narrative review. *Infectious Diseases and Therapy*, 14(5), 933–971. <https://doi.org/10.1007/s40121-025-01143-0>
- [7] Suleiman, T., Bello, M. A., Onyekwere, C. I., & Danjuma, S. (2025). Microbial health risks, antimicrobial contamination, and disease burden in Nigerian hospitals: A comprehensive meta-analysis [Preprint]. ResearchGate. <https://doi.org/10.13140/RG.2.2.40006.9993>
- [8] World Health Organization. (2022). Global report on infection prevention and control. World Health Organization. <https://iris.who.int/handle/10665/354489>
- [9] Onavbavba, G., Ogundele, A. O., Sanusi, M. B., & Adebayo, K. O. (2026). Knowledge, awareness, and perception of healthcare professionals towards Long COVID in Nigeria. *Frontiers in Public Health*, 14, Article 1768068. <https://doi.org/10.3389/fpubh.2026.1768068>
- [10] Oko, O. A., & Amali, J. A. (2026). Assessment of knowledge and practice of infection prevention and control among healthcare workers at Bauchi Specialist Hospital, Nigeria. *International Journal of Research Publication and Reviews*, 7(4), 7642–7652.
- [11] Massody, E. V., Okon, E. A., & Archibong, E. I. (2023). Knowledge, attitude and perceptions on infection prevention and control among primary healthcare workers in Calabar Municipality, Cross River State: A cross-sectional study. *International Journal of Health Psychology Research*, 3(3), 204–215.
- [12] Okonkwo, R., Obak, E., & Elonna, O. C. (2025). Knowledge, attitudes and practices of healthcare workers towards COVID-19 in three states of Nigeria in 2022. *BMJ Public Health*, 3(1), Article e001307. <https://doi.org/10.1136/bmjph-2024-001307>
- [13] Odetola, T. D., & Albert, S. O. (2026). Nurses' knowledge and perceived compliance with standard precautions in a tertiary health institution in south-western Nigeria. *Barnabas Journal of Nursing and Health Sciences*, 5(1), 14–25. <https://www.ajol.info/index.php/bjnhc/article/view/321726>

- [14] Ogoina, D., Mohammed, A., & Adekunle, O. J. (2025). Infection prevention practices and associated factors among clinical and non-clinical staff in two tertiary referral centers in southern Nigeria. *PLOS ONE*, 20(1), Article e0294810.
- [15] Okusiriki, F., Usman, H. O., & Bello, A. K. (2025). Skills gap and organizational performance in the healthcare sector of Nigeria: A conceptual review. *Governance and Research Journal of Business Management*, 4(2), 18–33.
<https://journals.amssr.org/grjbm/2025/12/12/skills-gap-and-organizational-performance-in-the-healthcare-sector-of-nigeria-a-conceptual-review/>
- [16] Yakubu, P. O., Aliyu, S. M., & Ibrahim, F. (2025). Evaluation of leadership competencies among public health managers in tertiary healthcare settings in Nigeria. *Journal of Health Management*, 27(2), 143–155.
- [17] Ogundele, A. O., Sanusi, M. B., & Adebayo, K. O. (2026). Knowledge, awareness, and perception of healthcare professionals towards Long COVID in Nigeria. *Frontiers in Public Health*, 14, 1768068.